antibodies .- online.com





anti-STAG2 antibody (AA 1021-1120)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL
Target:	STAG2
Binding Specificity:	AA 1021-1120
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This STAG2 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SA2
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	STAG2	
Alternative Name:	SA2 (STAG2 Products)	
Background:	Synonyms: Cohesin Subunit SA 2, Cohesin subunit SA-2, DKFZp686P168, DKFZp781H1753, SA	

2, SA2, SCC3 homolog 2, STAG 2, STAG 2, stag2, STAG2_HUMAN, Stromal antigen 2, FLJ25871, bA51701.1.

Background: Component of cohesin complex, a complex required for the cohesion of sister chromatids after DNA replication. The cohesin complex apparently forms a large proteinaceous ring within which sister chromatids can be trapped. At anaphase, the complex is cleaved and dissociates from chromatin, allowing sister chromatids to segregate. The cohesin complex may also play a role in spindle pole assembly during mitosis.

Gene ID: 10735

UniProt: Q8N3U4

Pathways: Stem Cell Maintenance

Application Details

Application Notes: WB 1:300-5000

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months