



[Go to Product page](#)

Datasheet for ABIN6989073  
**anti-TBPL1 antibody (AA 251-350)**

### Overview

Quantity:	100 µL
Target:	TBPL1
Binding Specificity:	AA 251-350
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TBPL1 antibody is un-conjugated
Application:	Western Blotting (WB)

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human TRF2
Isotype:	IgG
Cross-Reactivity:	Rat
Purification:	Purified by Protein A.

### Target Details

Target:	TBPL1
Alternative Name:	TRF2 ( <a href="#">TBPL1 Products</a> )
Background:	Synonyms: Telomeric DNA binding protein, Telomeric repeat binding factor 2, Telomeric repeat binding protein 2, TERF 2, TERF2, TRBF 2, TRBF2, TRF 2, TTAGGG repeat binding factor 2.

## Target Details

---

Background: Binds the telomeric double-stranded 5'-TTAGGG-3' repeat and plays a central role in telomere maintenance and protection against end-to-end fusion of chromosomes. In addition to its telomeric DNA-binding role, required to recruit a number of factors and enzymes required for telomere protection, including the shelterin complex, TERF2IP/RAP1 and DCLRE1B/Apollo. Component of the shelterin complex (telosome) that is involved in the regulation of telomere length and protection. Shelterin associates with arrays of double-stranded 5'-TTAGGG-3' repeats added by telomerase and protects chromosome ends, without its protective activity, telomeres are no longer hidden from the DNA damage surveillance and chromosome ends are inappropriately processed by DNA repair pathways. Together with DCLRE1B/Apollo, plays a key role in telomeric loop (T loop) formation by generating 3' single-stranded overhang at the leading end telomeres: T loops have been proposed to protect chromosome ends from degradation and repair. Required both to recruit DCLRE1B/Apollo to telomeres and activate the exonuclease activity of DCLRE1B/Apollo. Preferentially binds to positive supercoiled DNA. Together with DCLRE1B/Apollo, required to control the amount of DNA topoisomerase (TOP1, TOP2A and TOP2B) needed for telomere replication during fork passage and prevent aberrant telomere topology. Recruits TERF2IP/RAP1 to telomeres, thereby participating in to repressing homology-directed repair (HDR), which can affect telomere length.

---

Gene ID: 7014

UniProt: [Q15554](#)

## Application Details

---

Application Notes: WB 1:300-5000

Restrictions: For Research Use only

## Handling

---

Format: Liquid

Concentration: 1 µg/µL

Buffer: 0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: 4 °C, -20 °C

---

## Handling

---

Storage Comment: Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

---

Expiry Date: 12 months