antibodies - online.com







anti-VPS4B antibody (AA 301-400)



Overview

Quantity:	100 μL
Target:	VPS4B (vps4b)
Binding Specificity:	AA 301-400
Reactivity:	Human, Mouse, Fish
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This VPS4B antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human VPS4B
Isotype:	IgG
Cross-Reactivity:	Fish, Human, Mouse
Predicted Reactivity:	Rat,Cow,Pig
Purification:	Purified by Protein A.

Target Details

Target:	VPS4B (vps4b)		
---------	---------------	--	--

Target Details

Alternative Name:	VPS4B (vps4b Products)
Background:	Synonyms: 8030489C12Rik, Cell migration inducing 1, Cell migration-inducing gene 1 protein,
	MGC116271, MIG1, Protein SKD1, Skd1, SKD1B, Suppressor of K(+) transport growth defect 1,
	Suppressor of K+ transport defect 1, Vacuolar protein sorting 4 homolog B (S. cerevisiae),
	Vacuolar protein sorting 4 homolog B, Vacuolar protein sorting 4b, Vacuolar protein sorting-
	associated protein 4B, Vacuolar protein sorting-associating protein 4B, VPS4 2, VPS42, Vps4b,
	VPS4B_HUMAN.
	Background: Involved in late steps of the endosomal multivesicular bodies (MVB) pathway.
	Recognizes membrane-associated ESCRT-III assemblies and catalyzes their disassembly,
	possibly in combination with membrane fission. Redistributes the ESCRT-III components to the
	cytoplasm for further rounds of MVB sorting. MVBs contain intraluminal vesicles (ILVs) that are
	generated by invagination and scission from the limiting membrane of the endosome and
	mostly are delivered to lysosomes enabling degradation of membrane proteins, such as
	stimulated growth factor receptors, lysosomal enzymes and lipids. In conjunction with the
	ESCRT machinery also appears to function in topologically equivalent membrane fission
	events, such as the terminal stages of cytokinesis and enveloped virus budding (HIV-1 and
	other lentiviruses).
Gene ID:	9525
UniProt:	075351
Pathways:	Microtubule Dynamics, CXCR4-mediated Signaling Events
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months