



[Go to Product page](#)

Datasheet for ABIN6989203  
**anti-TOP2B antibody (AA 1-100)**

### Overview

Quantity:	100 µL
Target:	TOP2B
Binding Specificity:	AA 1-100
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TOP2B antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Topoisomerase II beta
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse,Dog,Cow,Sheep,Pig,Horse,Rabbit,Chimpanzee
Purification:	Purified by Protein A.

### Target Details

Target:	TOP2B
---------	-------

## Target Details

---

Alternative Name:	Topoisomerase II beta ( <a href="#">TOP2B Products</a> )
Background:	<p>Synonyms: Antigen MLAA 44, beta isozyme, DNA topoisomerase 2 beta, DNA topoisomerase 2-beta, DNA topoisomerase II 180 kD, DNA topoisomerase II, DNA topoisomerase II beta, DNA topoisomerase II beta isozyme, Top 2, TOP 2B, TOP IIB, Top2 beta, TOP2B, TOP2B_HUMAN, Top2beta, TOPIIB, Topo II beta, Topoisomerase (DNA) II beta 180 kDa, Topoisomerase (DNA) II beta, Topoisomerase IIb, U937 associated antigen.</p> <p>Background: Control of topological states of DNA by transient breakage and subsequent rejoining of DNA strands. Topoisomerase II makes double-strand breaks. Indirectly involved in vitamin D-coupled transcription regulation via its association with the WINAC complex, a chromatin-remodeling complex recruited by vitamin D receptor (VDR), which is required for the ligand-bound VDR-mediated transrepression of the CYP27B1 gene.</p>
Gene ID:	7155
UniProt:	<a href="#">Q02880</a>

## Application Details

---

Application Notes:	ELISA 1:500-1000 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 ICC 1:100-500
Restrictions:	For Research Use only

## Handling

---

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C

## Handling

---

Storage Comment: Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

---

Expiry Date: 12 months