

### Datasheet for ABIN6989257

# anti-PSMC3 antibody (AA 212-220)



Go to Product page

()	ve	r\/i	Δ	۱۸/
$\circ$	V C	1 V		v v

Quantity:	100 μL
Target:	PSMC3
Binding Specificity:	AA 212-220
Reactivity:	Human, Mouse, Xenopus laevis
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PSMC3 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human PSMC3
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Xenopus laevis
Predicted Reactivity:	Dog,Cow,Horse
Purification:	Purified by Protein A.

### **Target Details**

Target:	PSMC3			
---------	-------	--	--	--

## Target Details

Alternative Name:	PSMC3 (PSMC3 Products)	
Background:	Synonyms: 26S protease regulatory subunit 6A, 26S proteasome AAA-ATPase subunit RPT5,	
	Human immunodeficiency virus tat transactivator binding protein 1, MGC8487, Proteasome	
	(prosome macropain) 26S subunit ATPase 3, Proteasome 26S ATPase subunit 3, Proteasome	
	26S subunit ATPase 3, Proteasome subunit P50, PRS6A_HUMAN, PSMC 3, PSMC3, Tat binding	
	protein 1, TAT-binding protein 1, TBP-1, TBP1.	
	Background: The 26S protease is involved in the ATP-dependent degradation of ubiquitinated	
	proteins. The regulatory (or ATPase) complex confers ATP dependency and substrate	
	specificity to the 26S complex (By similarity). In case of HIV-1 infection, suppresses Tat-	
	mediated transactivation.	
Gene ID:	5702	
UniProt:	P17980	
Pathways:	Mitotic G1-G1/S Phases, DNA Replication, Synthesis of DNA, Ubiquitin Proteasome Pathway	
Application Details		
Application Notes:	WB 1:300-5000	
	ELISA 1:500-1000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
	ICC 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	

## Handling

Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months