

Datasheet for ABIN6989296

anti-Caspase 9 antibody (AA 231-330)



Overview

Quantity:	100 μL
Target:	Caspase 9 (CASP9)
Binding Specificity:	AA 231-330
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Caspase 9 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Caspase-9
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Cow,Sheep,Pig,Rabbit
Purification:	Purified by Protein A.

Target Details

Target: Caspase 9 (CASP9)

Target Details

Alternative Name:	Caspase-9 (CASP9 Products)
Background:	Synonyms: Caspase-9 subunit p35, Apaf-3, APAF 3, APAF3, Apoptosis related cysteine
	peptidase, Apoptotic protease activating factor 3, Apoptotic protease MCH 6, Apoptotic
	protease MCH6, CASP 9, CASP9, Caspase 9, Caspase 9 apoptosis related cysteine protease,
	Caspase 9 precursor, Caspase 9c, Caspase9, Caspase9 subunit p10, ICE LAP6, ICE like
	apoptotic protease 6, RNCASP9, MCH 6, MCH6, OTTHUMP00000044594, CASP9_HUMAN.
	Background: Caspase 9 (also known as ICE like apoptotic protease 6 (ICE LAP6), apoptotic
	protease Mch6, and apoptotic protease activating factor 3 (Apaf3)) is a member of the
	peptidase family C14 that contains a CARD domain. This caspase is active as a heterotetrame
	and has been reported to have two isoforms. ProCaspase 9 has been reported to be
	approximately 47 kD. This caspase is present in the cytosol and, upon activation, translocates
	to the mitochondria. Caspase 9 is involved in the caspase activation cascade responsible for
	apoptosis execution and cleaves/activates Caspase 3 and Caspase 6. Caspase 9 is inhibited by
	the dominant negative isoform, BcIXL, cIAP1, cIAP2, XIAP, and Livin. This caspase becomes
	activated when recruited to Apaf1/cytochrome c complex, and following cleavage by Apaf1,
	granzyme B, Caspase 3, possibly Caspase 8 and Caspase 10 into large p37 and small p10
	subunits. Caspase 9 intereacts with BIRC7 and has been shown to cleave PARP and vimentin.?
Gene ID:	842
UniProt:	P55211
Pathways:	MAPK Signaling, RTK Signaling, Apoptosis, Caspase Cascade in Apoptosis, Fc-epsilon Recepto
	Signaling Pathway, EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Positive
	Regulation of Endopeptidase Activity
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	FCM 1:20-100
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	ICC 1:100-500 For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months