# antibodies .- online.com







## anti-NEK6 antibody (AA 81-180)



$\sim$			
	N/6	1//r	$I \cap V$

Quantity:	100 μL
Target:	NEK6
Binding Specificity:	AA 81-180
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NEK6 antibody is un-conjugated
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human NEK6
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

### Target Details

### Target Details

Alternative Name:	NEK6 (NEK6 Products)	
Background:	Synonyms: Highly similar to cell cycle protein kinase CDC5/MSD2, NEK6, NEK6_HUMAN, Never	
	in mitosis A-related kinase 6, NIMA (Never In Mitosis Gene A) Related Kinase 6, NimA related	
	protein kinase 6, NimA-related protein kinase 6, Protein kinase SID6 1512, Protein kinase SID6-	
	1512, Putative serine threonine protein kinase, Serine/threonine protein kinase Nek6,	
	Serine/threonine-protein kinase Nek6, SID61512.	
	Background: Protein kinase which plays an important role in mitotic cell cycle progression.	
	Required for chromosome segregation at metaphase-anaphase transition, robust mitotic	
	spindle formation and cytokinesis. Phosphorylates ATF4, CIR1, PTN, RAD26L, RBBP6, RPS7,	
	RPS6KB1, TRIP4, STAT3 and histones H1 and H3. Phosphorylates KIF11 to promote mitotic	
	spindle formation. Involved in G2/M phase cell cycle arrest induced by DNA damage. Inhibition	
	of activity results in apoptosis. May contribute to tumorigenesis by suppressing p53/TP53-	
	induced cancer cell senescence.	
Gene ID:	10783	
UniProt:	Q9HC98	
Application Details		
Application Notes:	ELISA 1:500-1000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
	ICC 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	

### Handling

	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months