antibodies -online.com





anti-KIF13A antibody (AA 1501-1600)



Go to Product page

()	11/0	K\ /	iew
	\cup	'I V/I	$I \cap VV$

Quantity:	100 μL
Target:	KIF13A
Binding Specificity:	AA 1501-1600
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This KIF13A antibody is un-conjugated
Application:	ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffinembedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human KIF13A
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

|--|

Target Details

rarget Details		
Alternative Name:	KIF13A (KIF13A Products)	
Background:	Synonyms: bA500C11.2, FLJ27232, KI13A_HUMAN, KIF 13A, Kif13a, Kinesin family member	
	13A, Kinesin like protein KIF13A, Kinesin like protein RBKIN, Kinesin-like protein KIF13A, Kinesin-	
	like protein RBKIN, RBKIN.	
	Background: This gene encodes a member of the kinesin family of microtubule-based motor	
	proteins that function in the positioning of endosomes. This family member can direct	
	mannose-6-phosphate receptor-containing vesicles from the trans-Golgi network to the plasma	
	membrane, and it is necessary for the steady-state distribution of late endosomes/lysosomes.	
	It is also required for the translocation of FYVE-CENT and TTC19 from the centrosome to the	
	midbody during cytokinesis, and it plays a role in melanosome maturation. Alternative splicing	
	of this gene results in multiple transcript variants. [provided by RefSeq, Aug 2011]	
Gene ID:	63971	
UniProt:	Q9H1H9	
Application Details		
Application Notes:	ELISA 1:500-1000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
	ICC 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	

Handling

Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months