



[Go to Product page](#)

Datasheet for ABIN6989499 anti-MED28 antibody (AA 1-100)

Overview

Quantity:	100 µL
Target:	MED28
Binding Specificity:	AA 1-100
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MED28 antibody is un-conjugated
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human MED28
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit,Zebrafish
Purification:	Purified by Protein A.

Target Details

Target:	MED28
---------	-------

Target Details

Alternative Name:	MED28 (MED28 Products)
Background:	<p>Synonyms: EG1, Endothelial-derived protein 1, FKSG20, Magicin, med28, MED28_HUMAN, Mediator complex subunit 28, Mediator of RNA polymerase II transcription subunit 28, Mediator of RNA polymerase II transcription subunit 28, Mediator of RNA polymerase II transcription subunit 28, Merlin and Grb2-interacting cytoskeletal protein, Tumor angiogenesis marker EG-1.</p> <p>Background: Med28 is a 178 amino acid subunit of the Mediator complex that localizes to both nucleus and cytoplasm. Widely expressed with highest expression in vascular tissues such as placenta, testis and liver, Med28 participates in the regulation of Pol II-mediated gene expression and is thought to play a key role in signal regulation and cell proliferation. It is suggested that Med28 functions as a repressor of SMC (smooth muscle cell) differentiation and may be associated with abnormalities in SMC growth and differentiation, including atherosclerosis, asthma, hypertension and smooth muscle tumors.</p>
Gene ID:	610311
UniProt:	Q9H204
Pathways:	Stem Cell Maintenance , Regulation of Muscle Cell Differentiation

Application Details

Application Notes:	IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 ICC 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling

Storage:	4 °C,-20 °C
----------	-------------

Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
------------------	---

Expiry Date:	12 months
--------------	-----------