

Datasheet for ABIN6989500  
**anti-MAP2K7 antibody (pThr275)**



[Go to Product page](#)

## Overview

Quantity:	100 µL
Target:	MAP2K7
Binding Specificity:	pThr275
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MAP2K7 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin- embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))

## Product Details

Immunogen:	KLH conjugated synthesised phosphopeptide derived from human MEK7 around the phosphorylation site of Thr275
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Dog,Cow,Sheep,Pig,Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	MAP2K7
Alternative Name:	MEK7 ( <a href="#">MAP2K7 Products</a> )
Background:	<p>Synonyms: MEK7 (phospho T275), MEK7 (phospho S275), p-MEK7 (phospho S275), c-Jun N-terminal kinase kinase 2, Dual specificity mitogen activated protein kinase kinase 7, Dual specificity mitogen-activated protein kinase kinase 7, JNK activating kinase 2, JNK kinase 2, JNK-activating kinase 2, Jnkk 2, Jnkk-2, Jnkk2, MAP kinase kinase 7, MAP2K7, MAPK/ERK kinase 7, MAPKK 7, MAPKK-7, MAPKK7, MEK 7, Mitogen Activated Protein Kinase kinase 7, MKK 7, MKK-7, MKK7, MP2K7_HUMAN, PRKMK 7, PRKMK-7, PRKMK7, Sek 2, Sek-2, Sek2.</p> <p>Background: The protein encoded by this gene is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase specifically activates MAPK8/JNK1 and MAPK9/JNK2, and this kinase itself is phosphorylated and activated by MAP kinase kinase kinases including MAP3K1/MEKK1, MAP3K2/MEKK2, MAP3K3/MEKK5, and MAP4K2/GCK. This kinase is involved in the signal transduction mediating the cell responses to proinflammatory cytokines, and environmental stresses. Multiple alternatively spliced transcript variants encoding distinct isoforms have been found, but only one transcript variant has been supported and defined. [provided by RefSeq, Jul 2008].</p>
Gene ID:	5609
UniProt:	<a href="#">O14733</a>
Pathways:	<a href="#">MAPK Signaling</a> , <a href="#">TLR Signaling</a> , <a href="#">Fc-epsilon Receptor Signaling Pathway</a> , <a href="#">Activation of Innate immune Response</a> , <a href="#">Toll-Like Receptors Cascades</a> , <a href="#">BCR Signaling</a>

## Application Details

Application Notes:	WB 1:300-5000 ELISA 1:500-1000 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 ICC 1:100-500
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months