antibodies -online.com





Datasheet for ABIN6989505

anti-PRPSAP2 antibody (AA 1-100)



Go to Product page

/ N	11/0	K /	\sim	A /
	ve	1 \/	ı⊢۱	ΛI
\sim	' V C	1 V		v v

Quantity:	100 μL
Target:	PRPSAP2
Binding Specificity:	AA 1-100
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PRPSAP2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), ELISA, Immunocytochemistry (ICC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PRPSAP2
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	PRPSAP2		
---------	---------	--	--

Target Details

Alternative Name:	PRPSAP2 (PRPSAP2 Products)	
Background:	Synonyms: 41 kDa phosphoribosypyrophosphate synthetase associated protein, 41 kDa	
	phosphoribosypyrophosphate synthetase-associated protein, KPRB, KPRB_HUMAN,	
	MGC117304, MGC126719, MGC126721, PAP41, Phosphoribosyl pyrophosphate synthase-	
	associated protein 2, Phosphoribosyl pyrophosphate synthetase associated protein 2, PRPP	
	synthase-associated protein 2, PRPP synthetase associated protein 2, Prpsap2.	
	Background: This gene encodes a protein that associates with the enzyme	
	phosphoribosylpyrophosphate synthetase (PRS). PRS catalyzes the formation of	
	phosphoribosylpyrophosphate which is a substrate for synthesis of purine and pyrimidine	
	nucleotides, histidine, tryptophan and NAD. PRS exists as a complex with two catalytic subunits	
	and two associated subunits. This gene encodes a non-catalytic associated subunit of PRS.	
	Alternate splicing results in multiple transcript variants. [provided by RefSeq, Sep 2011]	
Gene ID:	5636	
UniProt:	060256	
Application Details		
Application Notes:	WB 1:300-5000	
	ELISA 1:500-1000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
	ICC 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	

Handling

	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months