antibodies -online.com





anti-PSMD14 antibody (AA 221-310)



Go to Product page

\sim				
	11/6	٦r١	/10	۱۸.

Quantity:	100 μL
Target:	PSMD14
Binding Specificity:	AA 221-310
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PSMD14 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PSMD14
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Cow,Sheep,Pig,Horse,Chicken,Rabbit,Zebrafish
Purification:	Purified by Protein A.

Target Details

Target: PSMD14

Target Details

Target Details		
Alternative Name:	PSMD14 (PSMD14 Products)	
Background:	Synonyms: 26S proteasome non-ATPase regulatory subunit 14, 26S proteasome regulatory	
	subunit rpn11, 26S proteasome-associated PAD1 homolog 1, 26S proteasome-associated	
	PAD1 homolog, PAD1, PAD1, yeast, homolog of, POH1, Proteasome (prosome, macropain) 26S	
	subunit, non-ATPase, 14, Proteasome 26S subunit non ATPase 14, PSDE_HUMAN, Psmd14,	
	RPN11.	
	Background: This gene encodes a component of the 26S proteasome. The 26S proteasome is a	
	large multiprotein complex that catalyzes the degradation of ubiquitinated intracellular proteins.	
	The encoded protein is a component of the 19S regulatory cap complex of the 26S proteasome	
	and mediates substrate deubiquitination. A pseudogene of this gene is also located on the long	
	arm of chromosome 2. [provided by RefSeq, Feb 2012]	
Gene ID:	10213	
UniProt:	000487	
Pathways:	Mitotic G1-G1/S Phases, DNA Replication, M Phase, Positive Regulation of Endopeptidase	
	Activity, Synthesis of DNA, Ubiquitin Proteasome Pathway	
Application Details		
· ·		
Application Notes:	WB 1:300-5000	
	ELISA 1:500-1000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
	ICC 1:100-500	
B		
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months