# antibodies - online.com







## anti-PDE3B antibody (AA 451-550)



$\sim$					
()	VE	۲۱	/1	$\triangle$	Λ

Quantity:	100 μL
	·
Target:	PDE3B
Binding Specificity:	AA 451-550
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PDE3B antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)),
	Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunocytochemistry (ICC),
	Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-
	embedded Sections) (IHC (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human PDE3B
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

#### Target Details

Target:	PDE3B		
---------	-------	--	--

#### **Target Details**

Alternative Name:	PDE3B (PDE3B Products)
Background:	Synonyms: 5"-cyclic phosphodiesterase B, Adipocyte cAMP phosphodiesterase cGMP inhibited,
	Al847709, cAMP Phosphodiesterase, adipocyte, cGMP-inhibited, CGI-PDE B, CGIP1, CGIPDE B,
	CGIPDE1, cGMP inhibited 3 5 cyclic phosphodiesterase B, CGMP inhibited cAMP
	phophodiesterase, cGMP-inhibited 3", Cyclic GMP inhibited phosphodiesterase B, Cyclic GMP-
	inhibited phosphodiesterase B, Cyclic nucleotide phosphodiesterase, HcGIP1, HcGIP1, rat,
	homolog of, Pde3b, PDE3B_HUMAN, Phosphodiesterase 3B cGMP inhibited.
	Background: PDE3B is found in lymphocytes. The PDE7A1 (HCP1) isozyme and the PDE7A2
	proteins, alternate splice products of PDE7A, are highly expressed in skeletal muscle. PDE7B is
	most highly expressed in pancreas. The PDE family contains proteins that serve tissue-specific
	roles in regulation of lipolysis, glycogenolysis, myocardial contractility, and smooth muscle
	relaxation.
Gene ID:	5140
UniProt:	Q13370
Pathways:	Negative Regulation of Hormone Secretion, Carbohydrate Homeostasis, cAMP Metabolic
	Process

Application Details		
Application Notes:	WB 1:300-5000	
	ELISA 1:500-1000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
	ICC 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	

### Handling

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months