## antibodies -online.com





## anti-SARS-CoV-2 NSP8 antibody (AA 40-90)



Go to Product page

| ( ) | 11/0   | K\ /   | iew                 | 1 |
|-----|--------|--------|---------------------|---|
|     | $\cup$ | 'I V/I | $\square \vee \vee$ | ı |
|     |        |        |                     |   |

| Overview             |   |  |
|----------------------|---|--|
| Quantity:            | 0.1 mg  |  |
| Target:              | SARS-CoV-2 NSP8 (NSP8)  |  |
| Binding Specificity: | AA 40-90  |  |
| Reactivity:          | SARS Coronavirus-2 (SARS-CoV-2)   |  |
| Host:                | Rabbit  |  |
| Clonality:           | Polyclonal  |  |
| Conjugate:           | This SARS-CoV-2 NSP8 antibody is un-conjugated  |  |
| Application:         | ELISA, Western Blotting (WB)  |  |
| Product Details      |   |  |
| Immunogen:           | Anti-SARS-CoV-2 (COVID-19) NSP8 antibody was raised against a peptide corresponding to 14   |  |
|                      | amino acids near the center of SARS-CoV-2 (COVID-19) NSP8 protein. The immunogen is         |  |
|                      | located within 40-90 amino acids of SARS-CoV-2 (COVID-19) NSP8.                             |  |
| Isotype:             | IgG   |  |
| Purification:        | SARS-CoV-2 (COVID-19) NSP8 Antibody is affinity chromatography purified via peptide column. |  |
|                      |   |  |
| Target Details       |   |  |
| Target:              | SARS-CoV-2 NSP8 (NSP8)  |  |
| Alternative Name:    | SARS-CoV-2 NSP8 (NSP8 Products)   |  |
| Target Type:         | Viral Protein   |  |

## Target Details

| Background:         | Coronavirus disease 2019 (COVID-19), formerly known as 2019-nCoV acute respiratory disease,     |  |
|---------------------|---|--|
|                     | is an infectious disease caused by SARS-CoV-2, a virus closely related to the SARS virus (1).   |  |
|                     | The disease is the cause of the 2019-20 coronavirus outbreak (2). The structure of 2019-nCoV    |  |
|                     | consists of the following: a spike protein (S), hemagglutinin-esterease dimer (HE), a membrane  |  |
|                     | glycoprotein (M), an envelope protein (E) a nucleoclapid protein (N) and RNA. NSP8 plays a role |  |
|                     | in viral RNA synthesis (3,4,5). Forms a hexadecamer with nsp7 (8 subunits of each) that may     |  |
|                     | participate in viral replication by acting as a primase. Alternatively, it may synthesize       |  |
|                     | substantially longer products than oligonucleotide primers (6).                                 |  |
| Gene ID:            | 43740578  |  |
| NCBI Accession:     | YP_009742615  |  |
| UniProt:            | PODTC1  |  |
| Application Details |   |  |
| Application Notes:  | WB: 0.5 μ,g/mL  |  |
|                     | Antibody validated: SARS-CoV-2 (COVID-19) NSP8 antibody can detect 2 ng of free peptide at 1    |  |
|                     | μ,g/mL in ELISA.  |  |
| Restrictions:       | For Research Use only   |  |
| Handling            |   |  |
| Format:             | Liquid  |  |
| Concentration:      | 1 mg/mL   |  |
| Buffer:             | SARS-CoV-2 (COVID-19) NSP8 Antibody is supplied in PBS containing 0.02 % sodium azide.          |  |
| Preservative:       | Sodium azide  |  |
| Precaution of Use:  | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which                   |  |
|                     | should be handled by trained staff only.  |  |
| Storage:            | -20 °C,4 °C   |  |
| Storage Comment:    | SARS-CoV-2 (COVID-19) NSP9 antibody can be stored at 4°C for three months and -20°C, stable     |  |
|                     | for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw   |  |
|                     | cycles. Antibodies should not be exposed to prolonged high temperatures.                        |  |