

Datasheet for ABIN6990285
anti-CIDEB antibody (C-Term)



[Go to Product page](#)

Overview

Quantity:	0.1 mg
Target:	CIDEB
Binding Specificity:	C-Term
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CIDEB antibody is un-conjugated
Application:	ELISA, Western Blotting (WB)

Product Details

Immunogen:	CIDE-B antibody was raised against a 16 amino acid peptide near the carboxy terminus of murine CIDE-B. The immunogen is located within the last 50 amino acids of CIDE-B.
Isotype:	IgG
Specificity:	It has no cross activity to CIDE-A.
Purification:	CIDE-B Antibody is affinity chromatography purified via peptide column.

Target Details

Target:	CIDEB
Alternative Name:	CIDE-B (CIDEB Products)
Background:	CIDE-B Antibody: Apoptosis is related to many diseases and induced by a family of cell death

Target Details

receptors and their ligands. Cell death signals are transduced by death domain containing adapter molecules and members of the caspase family of proteases. These death signals finally cause the degradation of chromosomal DNA by activated DNase. DFF45/ICAD has been identified as inhibitor of caspase activated DNase DFF40/CAD. DFF45 related proteins CIDE-A and CIDE-B (for cell death-inducing DFF-like effector A and B) were recently identified. CIDE contains a new type of domain termed CIDE-N, which has high homology with the regulatory domains of DFF45/ICAD and DFF40/CAD. Expression of CIDE-B induces apoptosis, which is inhibited by DFF45. CIDE-B is a DFF45-inhibitable effector that promotes cell death and DNA fragmentation. CIDE-B is expressed mainly in liver and at lower levels in spleen, kidney, peripheral blood lymphocytes, and bone marrow.

Molecular Weight: 25 kDa

Gene ID: 12684

UniProt: [O70303](#)

Application Details

Application Notes: CIDE-B antibody can be used for detection of CIDE-B by Western blot 0.5 µg/mL. An approximately 25 kDa band can be detected.

Antibody validated: Western Blot in mouse samples. All other applications and species not yet tested.

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 mg/mL

Buffer: CIDE-B Antibody is supplied in PBS containing 0.02 % sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C, 4 °C

Storage Comment: CIDE-B antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As

with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.