antibodies -online.com





Datasheet for ABIN6990444

anti-XAF1 antibody (C-Term)



()	1/0	r\ / I	014	
()	ve	I V I	-v	V

Quantity:	0.1 mg	
Target:	XAF1	
Binding Specificity:	C-Term	
Reactivity:	Human, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This XAF1 antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (IF)	
Product Details		
Immunogen:	XAF-1 antibody was raised against a synthetic peptide corresponding to amino acids at the C-terminus of human XAF-1. The immunogen is located within the last 50 amino acids of XAF-1.	
Isotype:	IgG	
Purification:	XAF-1 Antibody is affinity chromatography purified via peptide column.	
Target Details		
Target:	XAF1	
Alternative Name:	XAF-1 (XAF1 Products)	
Background:	XAF-1 Antibody: XAF-1 binds to XIAP, an inhibitor of caspases-3, -7, and -9, and triggers its relocation from the cytosol to the nucleus. Overexpression of XAF-1 results in the neutralization	

of XIAP's ability to inhibit cell death. XAF-1 is normally expressed in all adult and fetal tissues
but was found to be present in very low levels in a variety of cancer cell lines. In contrast, XIAP
levels have been shown to be high in a majority of cell lines. Low XAF-1 and high basal
expression of XIAP may therefore play a critical role in maintaining survival of cancer cell lines.
Both IFN-alpha2 and IFN-beta can induce XAF-1 mRNA in all cells examined but induction of
XAF-1 protein (as observed by immunoblot analysis) was seen only in cell lines sensitive to the
apoptotic effects of IFNs.

Molecular Weight:

Predicted: 35 kDa

Observed: 32 kDa

Gene ID:

54739

UniProt:

Q6GPH4

Application Details

Application Notes:

XAF-1 antibody can be used for the detection of XAF-1 by Western blot at 0.5 to 2 μ ,g/mL. Antibody can also be used for immunohistochemistry starting at 2 μ ,g/mL. For immunofluorescence start at 10 μ ,g/mL.

Antibody validated: Western Blot in human samples, Immunohistochemistry in human samples and Immunofluorescence in human samples. All other applications and species not yet tested.

Restrictions:

For Research Use only

Handling

Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	XAF-1 Antibody is supplied in PBS containing 0.02 % sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C,4 °C	
Storage Comment:	XAF-1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As	

with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.