antibodies -online.com





Overview

Target Details

Alternative Name:

DR4

DR4 (DR4 Products)

Target:

anti-DR4 antibody (N-Term)



Go to Product page

Quantity:	0.1 mg
Target:	DR4
Binding Specificity:	N-Term
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DR4 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC)
Product Details	
Immunogen:	Anti-DR4 antibody was raised against a peptide corresponding to 20 amino acids near the
	amino terminus of human DR4. The immunogen is located within the first 50 amino acids of
	DR4.
Isotype:	IgG
Specificity:	DR4 antibody has no cross reaction to DR5.
Purification:	DR4 Antibody is Antibody is DEAE purified.

Target Details

Background:	DR4 Antibody: Apoptosis, or programmed cell death, occurs during normal cellular
	differentiation and development of multicellular organisms. Apoptosis is induced by certain
	cytokines including TNF and Fas ligand in the TNF family through their death domain
	containing receptors, TNFR1 and Fas. A novel death domain containing receptor was recently
	identified and designated DR4 (for death receptor 4). The ligand for this novel death receptor
	has been identified and termed TRAIL, which is a new member in the TNF family. DR4 is also
	called TRAIL receptor-1 (TRAIL-R1). DR4 is expressed in most of human tissues including
	spleen, peripheral blood leukocytes, small intestine and thymus. Like TNFR1, Fas and DR3, DR4
	mediates apoptosis and NF-к,В activation in many tissues and cells.
Molecular Weight:	Predicted: 50kD
	Observed: 55kD (Post-modification: 1 N-linked glycosylation) kDa
Gene ID:	8797
UniProt:	000220
Pathways:	Apoptosis, Positive Regulation of Endopeptidase Activity
Application Details	
Application Notes:	WB: 1-4 μ,g/mL, ICC: 2 μ,g/mL, IF: 5 μ,g/mL.
	Antibody validated: Western Blot in human and mouse samples, Immunocytochemistry in
	human samples and Immunofluorescence in human and rat samples. All other applications
	and species not yet tested.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	DR4 Antibody is supplied in PBS containing 0.02 % sodium azide.
	DR4 Antibody is supplied in PBS containing 0.02 % sodium azide. Sodium azide

Handling

Storage:	-20 °C,4 °C
Storage Comment:	DR4 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As
	with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should
	not be exposed to prolonged high temperatures.