

Datasheet for ABIN6990607 **anti-AIF antibody (C-Term)**



[Go to Product page](#)

Overview

Quantity:	0.1 mg
Target:	AIF (AIFM1)
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB), ELISA, Immunocytochemistry (ICC)

Product Details

Immunogen:	Anti-AIF antibody was raised against a peptide corresponding to 14 amino acids near the carboxy terminus of human AIF. The immunogen is located within the last 50 amino acids of AIF.
Isotype:	IgG
Specificity:	Multiple isoforms of AIF are known to exist.
Purification:	AIF Antibody is affinity chromatography purified via peptide column.

Target Details

Target:	AIF (AIFM1)
Alternative Name:	AIF (AIFM1 Products)
Background:	AIF Antibody: Apoptosis is characterized by several morphological nuclear changes including chromatin condensation and nuclear fragmentation. These changes are triggered by the

Target Details

activation of members of caspase family, caspase activated DNase, and several novel proteins. A novel gene, the product of which causes chromatin condensation and DNA fragmentation, was recently identified, cloned, and designated apoptosis inducing factor (AIF). Like the critical molecules, cytochrome c and caspase-9, in apoptosis, AIF localizes in mitochondria. AIF translocates to the nucleus when apoptosis is induced and induces mitochondria to release the apoptogenic proteins cytochrome c and caspase-9. AIF induces chromatin condensation and DNA fragmentation, which are the hallmarks of apoptosis, of the isolated nucleus and the nucleus in live cells by microinjection. AIF is highly conserved between human and mouse and widely expressed.

Molecular Weight:	Predicted: 67kD
	Observed: 68kD kDa

Gene ID:	10256
----------	-------

UniProt:	O95381
----------	------------------------

Pathways:	Apoptosis , Positive Regulation of Endopeptidase Activity , Cell RedoxHomeostasis , Smooth Muscle Cell Migration , Warburg Effect
-----------	---

Application Details

Application Notes:	WB: 0.5-2 µg/mL, ICC: 5 µg/mL.
--------------------	--------------------------------

Antibody validated: Western Blot in human, mouse and rat samples, Immunocytochemistry in human samples. All other applications and species not yet tested.

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
---------	--------

Concentration:	1 mg/mL
----------------	---------

Buffer:	AIF Antibody is supplied in PBS containing 0.02 % sodium azide.
---------	---

Preservative:	Sodium azide
---------------	--------------

Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
--------------------	--

Handling

Storage:	-20 °C, 4 °C
----------	--------------

Storage Comment:	AIF antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
------------------	---