

Datasheet for ABIN6990647

anti-LY86 antibody (C-Term)



Overview

O V CI VIC VV	
Quantity:	0.1 mg
Target:	LY86
Binding Specificity:	C-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LY86 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	MD-1 antibody was raised against a 14 amino acid synthetic peptide from near the carboxy
	terminus of human MD-1. The immunogen is located within the last 50 amino acids of MD-1.
Isotype:	IgG
Purification:	
	MD-1 Antibody is affinity chromatography purified via peptide column.
Target Details	MD-1 Antibody is affinity chromatography purified via peptide column.
	MD-1 Antibody is affinity chromatography purified via peptide column. LY86
Target Details Target: Alternative Name:	
Target:	LY86
Target: Alternative Name:	LY86 MD-1 (LY86 Products)

rarget Details	
	TLRs is kept under tight control by the expression of endogenous inhibiting proteins such as RP105, a recently identified homolog of TLR4. This protein, in association with MD-1, interacts with and inhibits the TLR4/MD-2 signaling pathway. While MD-2 can directly bind to LPS, one of the activating molecules of TLR4, the function of MD-1 is less well-known. It has been suggested however, that the RP105/MD-1 complex influences antibody production mediated by both TLR4/MD-2 and TLR2 receptor complexes.
Gene ID:	9450
NCBI Accession:	NP_004262
UniProt:	095711
Pathways:	Toll-Like Receptors Cascades
Application Details	
Application Notes:	MD-1 antibody can be used for the detection of MD-1 by Western blot at 0.5 - 2 μ,g/mL.
	Antibody can also be used for immunohistochemistry starting at 2 μ ,g/mL.
	Antibody validated: Western Blot in human samples and Immunohistochemistry in human samples. All other applications and species not yet tested.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	MD-1 Antibody is supplied in PBS containing 0.02 % sodium azide.
Preservative:	Sodium azide

Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,4 °C
Storage Comment:	MD-1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.