

Datasheet for ABIN6990656
anti-BAG1 antibody (N-Term)



[Go to Product page](#)

Overview

Quantity:	0.1 mg
Target:	BAG1
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BAG1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	BAG-1 antibody was raised against a 14 amino acid synthetic peptide from near the amino terminus of human BAG-1. The immunogen is located within the first 50 amino acids of BAG-1.
Isotype:	IgG
Purification:	BAG-1 Antibody is affinity chromatography purified via peptide column.

Target Details

Target:	BAG1
Alternative Name:	BAG-1 (BAG1 Products)
Background:	BAG-1 Antibody: Bcl-2-associated athanogene 1 (BAG-1) was first identified as an anti-apoptotic bcl-2-binding protein. Later it was found to bind the molecular chaperones Hsp70 and Hsc70

Target Details

through its carboxy-terminal sequence (termed the Bag domain), resulting in the inhibition of the refolding activity of these chaperones. It is thought that by binding and inhibiting these molecular chaperones, BAG-1 is able to modulate the expression level of proteins requiring chaperones to fold correctly. One such group of proteins that are affected is glucocorticoid receptors. Other reports have suggested that the level of BAG-1 expression correlates with the aggressiveness of various cancers. Multiple isoforms of BAG-1 are known to exist.

Gene ID:	573
NCBI Accession:	NP_004314
UniProt:	Q99933

Application Details

Application Notes:	BAG-1 antibody can be used for the detection of BAG-1 by Western blot at 1 - 2 µg/mL. Antibody can also be used for immunohistochemistry starting at 2 µg/mL. For immunofluorescence start at 20 µg/mL. Antibody validated: Western Blot in human samples, Immunohistochemistry in human samples and Immunofluorescence in human samples. All other applications and species not yet tested.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	BAG-1 Antibody is supplied in PBS containing 0.02 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C, 4 °C
Storage Comment:	BAG-1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.