

Datasheet for ABIN6990695

anti-PRDM1 antibody (N-Term)



Go to Product page

_					
	1//	r	Vİ	\triangle	۸/
	V		VI		/ V

Quantity:	0.1 mg
Target:	PRDM1
Binding Specificity:	AA 100-150, N-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PRDM1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	Blimp-1 antibody was raised against a 17 amino acid synthetic peptide from near the amino
	terminus of human Blimp-1. The immunogen is located within amino acids 100 - 150 of Blimp-
	1.
Isotype:	IgG
Purification:	Blimp-1 Antibody is affinity chromatography purified via peptide column.
Target Details	
Target:	PRDM1
Alternative Name:	Blimp-1 (PRDM1 Products)
Background:	Blimp-1 Antibody: Blimp-1 was initially identified as a zinc finger-containing protein that drives
	the maturation of B lymphocytes into immunoglobulin-secreting cells. Together with X-box-

binding protein 1 (XBP1), Blimp-1 is induced upon terminal differentiation of plasma cells. The
transcriptional repressor activity of Blimp-1 has also been found to regulate T cell homeostasis
and function, possibly by suppressing the expression of the cytokines IL-2 and interferon-
gamma during T cell development. More recent experiments have suggested that Blimp-1 also
plays a major role in the formation of primordial germ cells (PGC) in developing mammalian
embryos. In these experiments, Blimp-1-deficient mutant mouse embryos form a cluster of
PGC-like cells which fail to show the expected migration, proliferation, and repression of
homeobox genes that normally accompany specification of primordial germ cells. Blimp-1
exists as at least two different isoforms.

Gene ID: 639

NCBI Accession: NP_001189

UniProt: 075626

Pathways: Regulation of Muscle Cell Differentiation

Application Details

Application Notes: Blimp-1 antibody can be used for detection of Blimp-1 by Western blot at 0.5 - 1 µ,g/mL.

Antibody can also be used for immunohistochemistry starting at 5 µ,g/mL.

Antibody validated: Western Blot in mouse samples and Immunohistochemistry in mouse samples. All other applications and species not yet tested.

Restrictions: For Research Use only

Handling

Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	Blimp-1 Antibody is supplied in PBS containing 0.02 % sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C,4 °C	
Storage Comment:	Blimp-1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As	

with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.