

Datasheet for ABIN6990756

anti-TBC1D1 antibody (C-Term)



Go to Product page

\sim		
	r\/I	$\triangle N$
\cup \vee \subset	1 V I	こい

ng
ID1
rm
an
oit
clonal
TBC1D1 antibody is un-conjugated
tern Blotting (WB), ELISA
1D1 antibody was raised against a 22 amino acid synthetic peptide from near the carboxy inus of human TBC1D1. The immunogen is located within the last 50 amino acids of 1D1.
1D1 Antibody is affinity chromatography purified via peptide column.
ID1
ID1 (TBC1D1 Products)
1D1 Antibody: TBC1D1 is the founding member of a family of proteins sharing a 180- to amino acid TBC domain and presumed to have a role in regulating cell growth and

differentiation. These proteins share significant homology with TRE2/USP6, yeast Bub2, and		
CDC16. TBC1D1 and TBC1D4 (AS160) have been demonstrated to be Rab GAPs (GT	Pase-	
activating proteins) that link upstream to Akt and phosphoinositide 3-kinase and dow	/nstream	
to Rabs involved in trafficking of GLUT4 vesicles. TBC1D1 regulates insulin-mediated GLUT4		
translocation through its GAP activity, and is a likely Akt substrate. Mutations in the Tbc1d1		
gene lead to some cases of severe human obesity.		

Molecular Weight:

Predicted: 128 kDa

Observed: 140 kDa

Gene ID:

23216

NCBI Accession:

NP_055988

UniProt:

Q86TI0

Application Details

Application Notes:

TBC1D1 antibody can be used for detection of TBC1D1 by Western blot at 2 μ,g/mL.

Antibody validated: Western Blot in human samples. All other applications and species not yet tested.

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	TBC1D1 Antibody is supplied in PBS containing 0.02 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,4 °C
Storage Comment:	TBC1D1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should

not be exposed to prolonged high temperatures.