

Datasheet for ABIN6990806

anti-DHX58 antibody (N-Term)



Overview

ar the amino
ar the amino 140 of LGP2.
140 of LGP2.

3	
	Toll-like receptors (TLR) 3, 7, 8, and 9 are expressed in immune cells and function as
	transmembrane pattern recognition receptors to detect foreign nucleic acids. Other proteins
	that play similar roles, such as RIG-1 and MDA5 are members of a CARD-helicase family and
	are expressed in the cytoplasm. A third protein, LGP2, is similar to RIG-1 and MDA5, except for
	lacking a homologous CARD domain. It is thought to act as an element of negative-feedback
	regulation of intracellular antiviral signaling. At least four isoforms of LGP2 are known to exist.
Gene ID:	79132
UniProt:	Q96C10
Application Details	

Analication	Matan
Application	Notes.

LGP2 antibody can be used for detection of LGP2 by Western blot at 0.5 - 1 μ ,g/mL.

Antibody validated: Western Blot in human samples. All other applications and species not yet tested.

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	LGP2 Antibody is supplied in PBS containing 0.02 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,4 °C
Storage Comment:	LGP2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.