

[Go to Product page](#)

Datasheet for ABIN6990824 **anti-GRIK4 antibody (N-Term)**

Overview

Quantity:	0.1 mg
Target:	GRIK4
Binding Specificity:	AA 30-80, N-Term
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GRIK4 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (IF)

Product Details

Immunogen:	Grik4 antibody was raised against a 14 amino acid synthetic peptide near the amino terminus of the human Grik4. The immunogen is located within amino acids 30 - 80 of Grik4.
Isotype:	IgG
Purification:	Grik4 Antibody is affinity chromatography purified via peptide column.

Target Details

Target:	GRIK4
Alternative Name:	Grik4 (GRIK4 Products)
Background:	Grik4 Antibody: Grik4 codes for the KA1 subunit of kainate-type ionotropic glutamate receptors which are critical regulators of network activity that act by modifying neuronal excitability,

Target Details

directly and indirectly, through GABAergic interneurons. Five subunits can assemble to form kainate receptors (KARs): GluR5 (coded by Grik1), GluR6 (coded by Grik2), and GluR7 (coded by Grik3) are the low-affinity subunits, and KA1 and KA2 are the high-affinity subunits. In the adult brain, KARs are located pre- and postsynaptically on pyramidal cells and on interneurons. Kainate receptors on GABA-containing interneurons enhance GABA release and thereby downregulate glutamatergic signaling. KARs have been implicated in numerous psychiatric disorders. Case control studies show significant association of Grik4 with both schizophrenia and bipolar disorder.

Gene ID:	2900
UniProt:	Q16099
Pathways:	Synaptic Membrane

Application Details

Application Notes:	<p>Grik4 antibody can be used for detection of Grik4 by Western blot at 0.5 - 2 µg/mL. Despite its predicted molecular weight, Grik4 often migrates at a lower molecular weight in SDS-PAGE. Antibody can also be used for immunohistochemistry starting at 2.5 µg/mL. For immunofluorescence start at 20 µg/mL.</p> <p>Antibody validated: Western Blot in rat samples, Immunohistochemistry in human samples and Immunofluorescence in human samples. All other applications and species not yet tested.</p>
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Grik4 Antibody is supplied in PBS containing 0.02 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C, 4 °C
Storage Comment:	Grik4 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As

with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.