

Datasheet for ABIN6990831

anti-SP110 antibody (N-Term)



| | | ۱۷(| e | ſ٧ | IE | W | V |
|--|--|-----|---|----|----|---|---|
|--|--|-----|---|----|----|---|---|

| Quantity: | 0.1 mg | |
|---|---|--|
| Target: | SP110 | |
| Binding Specificity: | AA 70-120, N-Term | |
| Reactivity: | Human, Mouse, Rat | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This SP110 antibody is un-conjugated | |
| Application: | Western Blotting (WB), ELISA | |
| Product Details | | |
| Immunogen: | IPR1 antibody was raised against a 16 amino acid synthetic peptide near the amino terminus of | |
| | the human IPR1. The immunogen is located within amino acids 70 - 120 of IPR1. | |
| | | |
| Isotype: | IgG | |
| Isotype: Purification: | IgG IPR1 Antibody is affinity chromatography purified via peptide column. | |
| | | |
| Purification: | | |
| Purification: Target Details | IPR1 Antibody is affinity chromatography purified via peptide column. | |
| Purification: Target Details Target: | IPR1 Antibody is affinity chromatography purified via peptide column. SP110 | |
| Purification: Target Details Target: Alternative Name: | IPR1 Antibody is affinity chromatography purified via peptide column. SP110 IPR1 (SP110 Products) | |
| Purification: Target Details Target: Alternative Name: | IPR1 Antibody is affinity chromatography purified via peptide column. SP110 IPR1 (SP110 Products) IPR1 Antibody: Susceptibility to tuberculosis (TB) in mice has recently been attributed to the | |

| transcription and may serve as a nuclear hormone receptor coactivator. Alternative splicing has |
|--|
| been observed for this gene and three transcript variants, encoding distinct isoforms, have been |
| identified. SP110 is the closest homolog of the IPR1 protein in humans. The IPR1/Sp110 gene |
| product might play a role in integrating signals generated by intracellular pathogens with |
| mechanisms controlling innate immunity, cell death, and pathogenesis. IPR1/Sp110 is up- |
| regulated after infection with M. tuberculosis and required by Anaplasma phagocytophilum for |
| infection of human promyelocytic cells. Defects in Sp110 are a cause of severely impaired |
| resistance to infection by M. tuberculosis. |

| Gene ID: | 3431 |
|-----------------|-----------|
| NCBI Accession: | NP_004501 |
| UniProt: | Q9HB58 |

Application Details

| Application Notes: | lpr1 antibody can be used for detection of lpr1 by Western blot at 1 - 2 μ,g/mL. |
|--------------------|---|
| | Antibody validated: Western Blot in human samples. All other applications and species not yet tested. |
| Restrictions: | For Research Use only |

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 mg/mL |
| Buffer: | IPR1 Antibody is supplied in PBS containing 0.02 % sodium azide. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | -20 °C,4 °C |
| Storage Comment: | IPR1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures. |