## antibodies .- online.com





## anti-BICD2 antibody (C-Term)



$\sim$	
( )\/△	rview
$\circ$	1 410 44

0.1 mg
BICD2
C-Term
Human, Mouse, Rat
Rabbit
Polyclonal
This BICD2 antibody is un-conjugated
Western Blotting (WB), ELISA
BICD2 antibody was raised against a 13 amino acid synthetic peptide from near the carboxy
BICD2 antibody was raised against a 13 amino acid synthetic peptide from near the carboxy terminus of human BICD2. The immunogen is located within the last 50 amino acids of BICD2.
terminus of human BICD2. The immunogen is located within the last 50 amino acids of BICD2.
terminus of human BICD2. The immunogen is located within the last 50 amino acids of BICD2.
terminus of human BICD2. The immunogen is located within the last 50 amino acids of BICD2.  IgG  This BICD2 antibody will not cross-react with BICD1.
terminus of human BICD2. The immunogen is located within the last 50 amino acids of BICD2.  IgG  This BICD2 antibody will not cross-react with BICD1.
terminus of human BICD2. The immunogen is located within the last 50 amino acids of BICD2.  IgG  This BICD2 antibody will not cross-react with BICD1.  BICD2 Antibody is affinity chromatography purified via peptide column.

protein that forms part of the cytoskeleton and mediates the correct sorting of mRNAs for
oocyte- and axis-determining factors during oogenesis. Similar to the highly homologous
protein BICD1, BICD2 can bind to dynein-dynactin complex, primarily through the dynamitin
subunit of dynactin. The C-terminus of BICD2 targets the protein to the Golgi complex while the
N-terminal domain of BICD2 co-immunoprecipitates with cytoplasmic dynein, suggesting
BICD2 plays a role in the dynein-dynactin interaction on the surface of membranous organelles.
Mice engineered to overexpress the BICD2 amino terminal domain in neurons developed
amyotrophic lateral sclerosis (ALS)-like features such as Golgi fragmentation, neurofilament
swelling in proximal axons, etc., suggesting that impaired dynein/dynactin function may explain
some of the pathological features observed in ALS patients.

Gene ID:	23299
UniProt:	Q8TD16
Pathways:	Maintenance of Protein Location

## **Application Details**

Application Notes:	BICD2 antibody can be used for detection of BICD2 by Western blot at 1 - 2 μ,g/mL.	
	Antibody validated: Western Blot in human samples. All other applications and species not yet tested.	
Restrictions:	For Research Use only	
Handling		

Tiaridility	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	BICD2 Antibody is supplied in PBS containing 0.02 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,4 °C
Storage Comment:	BICD2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should

not be exposed to prolonged high temperatures.