## antibodies -online.com







Go to Product page

$\sim$				
( )\	/arv	/1/	$\triangle I$	۸ /

Overview	
Quantity:	0.1 mg
Target:	SPRED1
Binding Specificity:	AA 120-170
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SPRED1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	Spred1 antibody was raised against a 20 amino acid synthetic peptide near the center of the
	human Spred1. The immunogen is located within amino acids 120 - 170 of Spred1.
Isotype:	IgG
Purification:	Spred1 Antibody is affinity chromatography purified via peptide column.
Target Details	
Target:	SPRED1
Alternative Name:	Spred1 (SPRED1 Products)
Background:	Spred1 Antibody: The Ras-MAP kinase pathway is essential for the differentiation of neuronal
	cells and myocytes, it is inhibited by Spred1, a member of the Sprouty family of proteins.
	Spred1 acts by suppressing the phosphorylation and activation of Raf. The Spred proteins have

n

Gene ID:	161742
NCBI Accession:	NP_689807
UniProt:	Q7Z699
Pathways:	Positive Regulation of Response to DNA Damage Stimulus

## **Application Details**

Application Notes:	Spred1 antibody can be used for detection of Spred1 by Western blot at 1 - 2 $\mu$ ,g/mL. Antibody
	can also be used for immunohistochemistry starting at 2.5 $\mu$ ,g/mL.

Antibody validated: Western Blot in mouse samples and Immunohistochemistry in human samples. All other applications and species not yet tested.

Restrictions: For Research Use only

## Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	Spred1 Antibody is supplied in PBS containing 0.02 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,4 °C
Storage Comment:	Spred1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should

not be exposed to prolonged high temperatures.