

Datasheet for ABIN6991324 anti-ANAPC4 antibody (N-Term)



Overview

Overview			
Quantity:	0.1 mg		
Target:	ANAPC4		
Binding Specificity:	AA 40-90, N-Term		
Reactivity:	Human, Mouse, Rat		
Host:	Rabbit		
Clonality:	Polyclonal		
Conjugate:	This ANAPC4 antibody is un-conjugated		
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))		
Product Details			
Immunogen:	APC4 antibody was raised against a 14 amino acid synthetic peptide near the amino terminus		
	of human APC4. The immunogen is located within amino acids 40 - 90 of APC4.		
Isotype:	IgG		
Purification:	APC4 Antibody is affinity chromatography purified via peptide column.		
Target Details			
Target:	ANAPC4		
Alternative Name:	APC4 (ANAPC4 Products)		
Background:	APC4 Antibody: Cell cycle regulated protein ubiquitination and degradation within subcellular		
	domains is thought to be essential for the normal progression of mitosis. APC4 is a highly		
	conserved component of the anaphase promoting complex/cyclosome (APC/C), a cell cycle-		

regulated E3 ubiquitin ligase that controls progression through mitosis and the G1 phase of the			
cell cycle. APC/C is responsible for degrading anaphase inhibitors, mitotic cyclins, and spindle-			
associated proteins ensuring that events of mitosis take place in proper sequence. The			
individual APC/C components mRNA and protein levels are expressed at approximately the			
same levels in most tissues and cell lines, suggesting that they perform their functions as part			
of a complex. While little is known of APC4, it is thought that APC4 associates with other APC/C			
components APC1, APC5, and CDC23 interdependently, such that loss of any one subunit			
reduces binding between the remaining three.			

Gene ID:	29945
UniProt:	Q9UJX5
Pathways:	M Phase

Application Details

Application Notes:	APC4 antibody can be used for detection of APC4 by Western blot at 1 - 2 µ,g/mL. Antibody can	
	also be used for immunohistochemistry starting at 5 μ ,g/mL.	

Antibody validated: Western Blot in mouse samples and Immunohistochemistry in rat samples. All other applications and species not yet tested.

Restrictions:	For Research Use only

Handling

Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	APC4 Antibody is supplied in PBS containing 0.02 % sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C,4 °C	
Storage Comment:	APC4 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.	