antibodies - online.com





anti-Secernin 1 antibody (C-Term)



Coto	Product	nage
GU 10	FIUUUUU	paye

()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	0.1 mg
Target:	Secernin 1 (SCRN1)
Binding Specificity:	AA 320-370, C-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Secernin 1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (IF)
Product Details	
Immunogen:	SCRN1 antibody was raised against a 20 amino acid synthetic peptide near the carboxy terminus of human SCRN1. The immunogen is located within amino acids 320 - 370 of SCRN1.
Isotype:	IgG
Purification:	SCRN1 Antibody is affinity chromatography purified via peptide column.
Target Details	
Target:	Secernin 1 (SCRN1)
Alternative Name:	SCRN1 (SCRN1 Products)
Background:	SCRN1 Antibody: SCRN1 was first identified as a cytosolic protein that is involved in the regulation of exocytosis from peritoneal mast cells. More recent studies have shown that

Gene ID:	9805
NCBI Accession:	NP_001138985
UniProt:	012765

Application Details

Application Notes:	SCRN1 antibody can be used for detection of SCRN	N1 by Western blot at 1 μ,g/mL. Antibody can
--------------------	--	--

also be used for immunohistochemistry starting at 5 $\mu\text{,g/mL}.$ For immunofluorescence start at

20 μ,g/mL.

Antibody validated: Western Blot in human samples, Immunohistochemistry in human samples

and Immunofluorescence in human samples. All other applications and species not yet tested.

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	SCRN1 Antibody is supplied in PBS containing 0.02 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,4 °C
Storage Comment:	SCRN1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.