

Datasheet for ABIN6991385

anti-WDR35 antibody (N-Term)



Overview

| Quantity: | 0.1 mg |
|----------------------|---|
| Target: | WDR35 |
| Binding Specificity: | AA 60-110, N-Term |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This WDR35 antibody is un-conjugated |
| Application: | ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |
| Product Details | |
| Immunogen: | WDR35 antibody was raised against a 16 amino acid synthetic peptide near the amino terminus |
| | of human WDR35. The immunogen is located within amino acids 60 - 110 of WDR35. |
| Isotype: | IgG |
| Specificity: | WDR35 antibody is human specific. |
| Purification: | WDR35 Antibody is affinity chromatography purified via peptide column. |
| Target Details | |
| Target: | WDR35 |
| Alternative Name: | WDR35 (WDR35 Products) |
| Background: | WDR35 Antibody: WD40 repeats are a common structural module in eukaryotic proteins, and |
| | |

| proteins containing WD40 domains have a wide range of functions, including signal |
|--|
| transduction, cell cycle regulation, RNA splicing, and transcription. One such protein, WDR35, |
| also known as CED2, has been shown to be mutated in patients with Sensenbrenner |
| syndrome/cranioectodermal dysplasia (CED), an autosomal-recessive disease that is |
| characterized by craniosynstosis and ectodermal and skeletal abnormalities. WDR35 localizes |
| to cilia and dentrosomes during embryogenesis and human and mouse fibroblasts that lack |
| this gene fail to produce cilia. Mutations in this gene can also cause short-rib polydactyly |
| syndromes due to abnormal ciliogenesis. |

| Gene ID: | 57539 |
|-----------------|--------------|
| NCBI Accession: | NP_001006658 |
| UniProt: | Q9P2L0 |
| | |

Pathways: Hedgehog Signaling

Application Details

| Application Notes: | WDR35 antibody can be used for detection of WDR35 by immunohistochemistry at 5 μ ,g/mL. |
|--------------------|---|
| | Antibody validated: Immunohistochemistry in human samples. All other applications and |
| | species not yet tested. |
| Restrictions: | For Research Use only |

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 mg/mL |
| Buffer: | WDR35 Antibody is supplied in PBS containing 0.02 % sodium azide. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | -20 °C,4 °C |
| Storage Comment: | WDR35 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should |

not be exposed to prolonged high temperatures.