

## Datasheet for ABIN6991393

## anti-YBX2 antibody (AA 300-350)



Go to Product page

| $\sim$ |             |     |     |   |
|--------|-------------|-----|-----|---|
| $O_1$  | <b>/</b> el | rVI | 161 | Λ |

| 0.1 mg<br>YBX2   |
|--|
| YBX2   |
|  |
| AA 300-350   |
| Human, Mouse, Rat  |
| Rabbit   |
| Polyclonal   |
| This YBX2 antibody is un-conjugated  |
| Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunohistochemistry (Paraffin-         |
| embedded Sections) (IHC (p))   |
|  |
| YBX2 antibody was raised against an 18 amino acid synthetic peptide near the carboxy end of    |
| human YBX2. The immunogen is located within amino acids 300 - 350 of YBX2.                     |
| IgG  |
| YBX2 Antibody is affinity chromatography purified via peptide column.                          |
|  |
| YBX2   |
| YBX2 (YBX2 Products)   |
|  |
| YBX2 Antibody: Germ cell specific Y-box binding protein (YBX2), a germ-cell-specific member of |
|  |

| ribonucleoprotein particles (mRNPs) and involved in the regulation of the stability and/or      |
|---|
| translation of germ cell mRNAs. It is proposed to function as a coactivator of transcription in |
| the nucleus and to stabilize and store maternal and paternal mRNAs in the cytoplasm. YBX2       |
| binds to the Y-box consensus promoter element and is expressed in oocytes and testicular        |
| germ cells in the stage of spermatogonia to spermatocyte. Recent studies show that deletion     |
| of YBX2 leads to the disruption of both spermatogenesis and oogenesis.                          |
|   |

| Gene ID:        | 51087             |
|-----------------|-------------------|
| NCBI Accession: | NP_057066         |
| UniProt:        | Q9Y2T7            |
| Pathways:       | Chromatin Binding |

## **Application Details**

Application Notes: YBX2 antibody can be used for detection of YBX2 by Western blot at 1 - 2  $\mu$ ,g/mL. Antibody can also be used for immunohistochemistry starting at 10  $\mu$ ,g/mL. For immunofluorescence start at

 $20~\mu,g/mL$ .

Antibody validated: Western Blot in human samples, Immunohistochemistry in human samples and Immunofluorescence in human samples. All other applications and species not yet tested.

Restrictions: For Research Use only

## Handling

| Format:            | Liquid   |
|--------------------|--|
| Concentration:     | 1 mg/mL  |
| Buffer:            | YBX2 Antibody is supplied in PBS containing 0.02 % sodium azide.   |
| Preservative:      | Sodium azide   |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.   |
| Storage:           | -20 °C,4 °C  |
| Storage Comment:   | YBX2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures. |