



[Go to Product page](#)

Datasheet for ABIN6991428
anti-PLAC2 antibody (AA 60-110)

Overview

Quantity:	0.1 mg
Target:	PLAC2 (TINCR)
Binding Specificity:	AA 60-110
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PLAC2 antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunocytochemistry (ICC), Immunofluorescence (IF)

Product Details

Immunogen:	PLAC2 antibody was raised against a 17 amino acid synthetic peptide near the center of human PLAC2. The immunogen is located within amino acids 60 - 110 of PLAC2.
Isotype:	IgG
Purification:	PLAC2 Antibody is affinity chromatography purified via peptide column.

Target Details

Target:	PLAC2 (TINCR)
Alternative Name:	PLAC2 (TINCR Products)
Background:	PLAC2 Antibody: PLAC2 (placenta-specific protein 2) is a 151 amino acid protein that is encoded by a gene located on human chromosome 19p13. Based on its amino acid sequence, PLAC2 is predicted to be a multi-pass membrane protein. There are no reported studies or

Target Details

functions reported for this protein.

Gene ID: 257000

UniProt: [Q9NPR7](#)

Application Details

Application Notes: PLAC2 antibody can be used for detection of PLAC2 by Western blot at 1 - 2 μ g/mL. Antibody can also be used for immunocytochemistry starting at 10 μ g/mL. For immunofluorescence start at 20 μ g/mL.

Antibody validated: Western Blot in human samples, Immunocytochemistry in human samples and Immunofluorescence in human samples. All other applications and species not yet tested.

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 mg/mL

Buffer: PLAC2 Antibody is supplied in PBS containing 0.02 % sodium azide.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C, 4 °C

Storage Comment: PLAC2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.