antibodies - online.com





anti-TOX antibody (N-Term)



Overview	
Quantity:	0.1 mg
Target:	TOX
Binding Specificity:	AA 130-180, N-Term
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TOX antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF)
Product Details	
Immunogen:	TOX antibody was raised against a 17 amino acid synthetic peptide near the amino terminus of
	human TOX. The immunogen is located within amino acids 130 - 180 of TOX.
Isotype:	IgG
Specificity:	No alternatively spliced transcript variants have been observed.
Purification:	TOX Antibody is affinity chromatography purified via peptide column.
Target Details	
Target:	TOX
Alternative Name:	TOX (TOX Products)
Background:	TOX Antibody: TOX (thymocyte selection-associated high mobility group (HMG) box protein) is

Molecular Weight:	58 kDa
	development of class II major histocompatibility complex-specific T cells.
	calcineurin dependent. TOX-dependent transition to the CD4+CD8 stage is required for
	thymocytes. TOX-mediated positive selection is associated with up-regulation of RUNX3 and is
	TOX is upregulated by pre-T cell receptor (pre-TCR) and TCR activation in immature
	regulation of T-cell development. It is a 526 amino acid nuclear protein and the expression of
	a member of the HMG box family of DNA-binding proteins and likely plays a role in the

Molecular Weight:	58 kDa
Gene ID:	9760
NCBI Accession:	NP_055544
UniProt:	094900

Application Details

Application Notes:	TOX antibody can be used for detection of TOX by Western blot at 1 - 2 μ ,g/mL. For

immunofluorescence start at 20 μ ,g/mL.

Antibody validated: Western Blot in human samples and Immunofluorescence in human samples. All other applications and species not yet tested.

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	TOX Antibody is supplied in PBS containing 0.02 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,4 °C
Storage Comment:	TOX antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.