antibodies -online.com





anti-MTERFD2 antibody (C-Term)



Go to	Product	page

_							
0	V	е	r١	/1	е	V	1

Quantity:	0.1 mg
Target:	MTERFD2
Binding Specificity:	AA 260-310, C-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MTERFD2 antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunofluorescence (IF), Immunohistochemistry (Paraffin-
	embedded Sections) (IHC (p))
Product Details	
Jagot Detailo	
Immunogen:	MTERFD2 antibody was raised against a 17 amino acid peptide near the carboxy terminus of
	MTERFD2 antibody was raised against a 17 amino acid peptide near the carboxy terminus of human MTERFD2. The immunogen is located within amino acids 260 - 310 of MTERFD2.
Immunogen:	human MTERFD2 . The immunogen is located within amino acids 260 - 310 of MTERFD2.
Immunogen: Isotype:	human MTERFD2 . The immunogen is located within amino acids 260 - 310 of MTERFD2. IgG
Immunogen: Isotype:	human MTERFD2 . The immunogen is located within amino acids 260 - 310 of MTERFD2. IgG At least three isoforms of MTERFD2 are known to exist. MTERFD2 antibody is predicted to not
Immunogen: Isotype: Specificity:	human MTERFD2 . The immunogen is located within amino acids 260 - 310 of MTERFD2. IgG At least three isoforms of MTERFD2 are known to exist. MTERFD2 antibody is predicted to not cross-react with other MTERFD protein family members.
Immunogen: Isotype: Specificity: Purification:	human MTERFD2 . The immunogen is located within amino acids 260 - 310 of MTERFD2. IgG At least three isoforms of MTERFD2 are known to exist. MTERFD2 antibody is predicted to not cross-react with other MTERFD protein family members.

Target Details

Background:	MTERFD2 Antibody: Members of the mTERF (mitochondrial transcription termination factor)
	family, are mitochondrial proteins that are believed to be transcription termination factors.
	MTERFD2 is targeted to the mitochondria and is ubiquitously expressed, with highest
	expression levels in fore- and midbrain, diencephalon, spinal cord, tongue, lung liver and kidney.
	MTERFD2 has been suggested to play a role in organ differentiation during embryogenesis. A
	closely related mTERF family member, MTERFD3, is believed to be involved in cell cycle
	regulation and cell growth by modulating mitochondrial transcription.
Molecular Weight:	Predicted: 42 kDa
	Observed: 44 kDa
Gene ID:	130916
NCBI Accession:	NP_872307
UniProt:	Q7Z6M4
Application Details	
Application Notes:	MTERFD2 Antibody can be used for detection of MTERFD2 by Western blot at 1 μ,g/mL.
	Antibody validated: Western Blot in human samples, Immunohistochemistry in human samples
	and Immunofluorescence in human samples. All other applications and species not yet tested.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	MTERFD2 Antibody is supplied in PBS containing 0.02 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	-20 °C,4 °C
Storage Comment:	MTERFD2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.