antibodies -online.com







anti-IFIT1 antibody (C-Term)



()	11/	IN	/ie	A .
	/ // 	۱ ات	/ (−	' \/\/

Overview		
Quantity:	0.1 mg	
Target:	IFIT1	
Binding Specificity:	AA 290-340, C-Term	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This IFIT1 antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF), Immunohistochemistry (Paraffinembedded Sections) (IHC (p))	
Product Details		
Immunogen:	IFIT1 antibody was raised against an 18 amino acid peptide near the carboxy terminus of human IFIT1. The immunogen is located within amino acids 290 - 340 of IFIT1.	
Isotype:	IgG	
Specificity:	IFIT1 antibody is human, mouse and rat reactive. At least three isoforms of IFIT1 are known to exist. This antibody is predicted to not cross-react with other members of the IFIT protein family.	
Purification:	IFIT1 antibody is affinity chromatography purified via peptide column.	
Target Details		
Target:	IFIT1	

Target Details

Alternative Name:	IFIT1 (IFIT1 Products)	
Background:	The interferon-induced protein with tetratricopeptide repeats 1 (IFIT1) protein is a member of a family of tetratricopeptide repeat-containing proteins whose transcription is upregulated by interferons, virus infection, and molecular patterns such as dsRNA or lipopolysaccharides (1,2). These proteins have been suggested to induce anti-viral cellular activities in response to infection (2). Together with the interferon-induced transmembrane protein 1 (IFITM1), IFIT1 inhibits the replication of the Hepatitis C virus, suggesting that it may be useful in the development of therapeutic treatments (3).	
Molecular Weight:	Predicted: 53 kDa	
	Observed: 53 kDa	
Gene ID:	3434	
NCBI Accession:	NP_001539	
UniProt:	P09914	
Pathways:	Hepatitis C	
Application Details		
Application Notes:	IFIT1 antibody can be used for detection of IFIT1 by Western blot at 1 - 2 μ ,g/mL. Antibody can also be used for Immunohistochemistry at 5 μ ,g/mL. For Immunoflorescence start at 20 μ , g/mL.	
	Antibody validated: Western Blot in rat samples, Immunohistochemistry in human samples and Immunofluorescence in human samples. All other applications and species not yet tested.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	IFIT1 Antibody is supplied in PBS containing 0.02 % sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	

Handling

	should be handled by trained staff only.	
Storage:	-20 °C,4 °C	
Storage Comment:	IFIT1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.	