

## Datasheet for ABIN6992077

## anti-IQSEC2 antibody (C-Term)



## Overview

Quantity:	0.1 mg
Target:	IQSEC2
Binding Specificity:	AA 1380-1430, C-Term
Reactivity:	Mouse, Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IQSEC2 antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Immunogen:	IQSEC2 antibody was raised against an 18 amino acid peptide near the carboxy terminus of
	human IQSEC2. The immunogen is located within amino acids 1380 - 1430 of IQSEC2.
Isotype:	IgG
Specificity:	IQSEC2 antibody is human, mouse and rat reactive. At least two isoforms of IQSEC2 are known
	to exist, this antibody will only detect the larger isoform. IQSEC2 antibody is predicted to not
	cross-react with IQSEC1.
Purification:	IQSEC2 antibody is affinity chromatography purified via peptide column.
Target Details	
Target:	IQSEC2
Alternative Name:	IQSEC2 (IQSEC2 Products)

## **Target Details**

901 = 0000	
Background:	ADP ribosylation factors (Arfs) are small GTP-binding proteins known for their role in vesicular
	transport, where they nucleate the assembly of coat protein complexes at sites of carrier
	vesicle formation. Similar to the related protein IQSEC1, IQSEC2 is a guanine nucleotide
	exchange protein for ARF1 and ARF6 that belongs to the BRAG family and contains one IQ
	domain, one PH domain and one SEC7 domain (1,2). IQSEC2 is a major component of the
	postsynaptic density (PSD) and colocalizes with PSD-95. It has been suggested to modulate the
	functions of the ARF proteins (1). Mutations in this gene have been implicated in nonsyndromic
	X-linked mental retardation (3).
Molecular Weight:	Predicted: 104, 133, 141, 164 kDa
	Observed: 140 kDa
Gene ID:	23096
NCBI Accession:	NP_001104595
UniProt:	Q5JU85
Application Details	
Application Notes:	IQSEC2 antibody can be used for detection of IQSEC2 by Western blot at 1 - 2 μ,g/mL. Antibody
	can also be used for immunohistochemistry starting at 5 $\mu$ ,g/mL.
	Antibody validated: Western Blot in human samples and Immunohistochemistry in mouse
	samples. All other applications and species not yet tested.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	IQSEC2 antibody is supplied in PBS containing 0.02 % sodium azide.
	IQSEC2 antibody is supplied in PBS containing 0.02 % sodium azide.  Sodium azide
Preservative:	
Buffer: Preservative: Precaution of Use:	Sodium azide

Storage C	Comment:
oto.ago c	

IQSEC2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.