

Datasheet for ABIN6992152 **anti-LSD1 antibody (N-Term)**

[Go to Product page](#)

Overview

Quantity:	0.1 mg
Target:	LSD1 (KDM1A)
Binding Specificity:	AA 90-140, N-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LSD1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	Anti-LSD1 antibody was raised against a peptide corresponding to 19 amino acids near the amino terminus of human LSD1. The immunogen is located within amino acids 90-140 of LSD1.
Isotype:	IgG
Specificity:	At least two isoforms of LSD1 are known to exist, this antibody detects both.
Purification:	LSD1 antibody is affinity chromatography purified via peptide column.

Target Details

Target:	LSD1 (KDM1A)
Alternative Name:	LSD1 (KDM1A Products)

Target Details

Background:	Histone modifications mediate changes in gene expression by altering chromatin structure or by serving as a platform to recruit other proteins. LSD1 is a recently discovered amine oxidase that catalyzes the lysine-specific demethylation of histone proteins via an FAD-dependent oxidative reaction (1). Methylation on histone H3-K9 is thought to play an important role in heterochromatin formation, while methylation on arginine and some lysine residues (such as H3-K4) is associated with active transcription (2). LSD1 associates with various proteins, including HDAC1/2, CoREST, and BHC80, that act to regulate LSD1 activity in vivo, and in a histone H3-K4-specific methylase complex that is involved in transcriptional regulation (3,4). Experiments have shown that CoREST, a SANT domain-containing corepressor (5) acts to enhance LSD1 activity, while BHC80, a PHD domain-containing protein (6), inhibits CoREST/LSD1 activity in vitro (3). LSD1-mediated histone demethylation thus may have significant effects on gene expression.
Molecular Weight:	Predicted: 93kD Observed: 93 kD kDa
Gene ID:	23028
NCBI Accession:	NP_001009999
UniProt:	O60341
Pathways:	Regulation of Hormone Metabolic Process , Regulation of Hormone Biosynthetic Process , Negative Regulation of intrinsic apoptotic Signaling , Warburg Effect

Application Details

Application Notes:	WB: 1 - 2 µg/mL, IHC-P: 5 µg/mL. Antibody validated: Western Blot in human, mouse and rat samples, Immunohistochemistry in human samples. All other applications and species not yet tested.
--------------------	--

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	LSD1 antibody is supplied in PBS containing 0.02 % sodium azide.

Handling

Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C, 4 °C
Storage Comment:	LSD1 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.