

Datasheet for ABIN6992168

anti-DIS3L2 antibody (C-Term)



Overview

Quantity:	0.1 mg
Target:	DIS3L2
Binding Specificity:	AA 720-770, C-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DIS3L2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA
Product Details	
Immunogen:	DIS3L2 antibody was raised against a 17 amino acid peptide near the carboxy terminus of
	human DIS3L2. The immunogen is located within amino acids 720 - 770 of DIS3L2.
Isotype:	IgG
Specificity:	
Specificity:	DIS3L2 antibody is human, mouse and rat reactive. At least three isoforms of DIS3L2 are known
Specificity:	DIS3L2 antibody is human, mouse and rat reactive. At least three isoforms of DIS3L2 are known to exist, this antibody will only detect the longest isoform. DIS3L2 is predicted to not cross-react
Specificity :	
Specificity: Purification:	to exist, this antibody will only detect the longest isoform. DIS3L2 is predicted to not cross-react
	to exist, this antibody will only detect the longest isoform. DIS3L2 is predicted to not cross-react with DIS3 or DIS3L.
Purification:	to exist, this antibody will only detect the longest isoform. DIS3L2 is predicted to not cross-react with DIS3 or DIS3L.
Purification: Target Details	to exist, this antibody will only detect the longest isoform. DIS3L2 is predicted to not cross-react with DIS3 or DIS3L. DIS3L2 antibody is affinity chromatography purified via peptide column.

Target Details

Background:	The exosome is involved in a multitude of cellular RNA processing and degradation events (1). DIS3, also known as exosome complex exonuclease RRP44, is a ribonuclease that acts directly in the processing, turnover, and surveillance of a large number of distinct RNA species (2). DIS3L2 is a paralog of DIS3, but unlike DIS3 is located in the cytoplasm (3). DIS3L2 defines a novel eukaryotic RNA degradation pathway and preferentially targets uridylated RNA (4). DIS3L2 mutations have also been associated with sporadic Wilms tumors (3).
Molecular Weight:	Predicted: 97 kDa
	Observed: 94 kDa
Gene ID:	129563
NCBI Accession:	NP_689596
UniProt:	Q8IYB7
Pathways:	Stem Cell Maintenance
Application Details	
Application Notes:	DIS3L2 antibody can be used for detection of DIS3L2 by Western blot at 1 - 2 μ ,g/mL. Antibody can also be used for immunohistochemistry starting at 5 μ ,g/mL.
	Antibody validated: Western Blot in mouse samples and Immunohistochemistry in rat samples. All other applications and species not yet tested.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	DIS3L2 antibody is supplied in PBS containing 0.02 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C,4 °C
Storage Comment:	DIS3L2 antibody can be stored at 4°C for three months and -20°C, stable for up to one year.