



[Go to Product page](#)

Datasheet for ABIN6992404

SARS-CoV-2 Spike S1 Protein (B.1.351 - beta) (Fc Tag)

2 Images

Overview

Quantity:	100 µg
Target:	SARS-CoV-2 Spike S1
Protein Characteristics:	B.1.351 - beta
Origin:	SARS Coronavirus-2 (SARS-CoV-2), SARS CoV-2 Beta
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This SARS-CoV-2 Spike S1 protein is labelled with Fc Tag.

Product Details

Purpose:	SARS-CoV-2 S1 protein (L18F, D80A, D215G, LAL242-244del, R246I, K417N, E484K, N501Y, D614G), Fc Tag
Sequence:	AA 16-685
Characteristics:	SARS-CoV-2 S1 protein, Fc Tag is expressed from human 293 cells (HEK293). It contains AA Val 16 - Arg 685 (Accession # QHD43416.1). The mutations (L18F, D80A, D215G, LAL242-244del, R246I, K417N, E484K, N501Y, D614G) were identified in the SARS-CoV-2 variant (known as B.1.351 or 20C/501Y.V2) which emerged in the South Africa.
Purity:	>95 % as determined by SDS-PAGE.
Endotoxin Level:	Less than 1.0 EU per µg by the LAL method.

Target Details

Target: SARS-CoV-2 Spike S1

Alternative Name: SARS-CoV-2 S1 protein ([SARS-CoV-2 Spike S1 Products](#))

Target Type: Viral Protein

Background: It's been reported that Coronavirus can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.

Molecular Weight: 101.1 kDa

Application Details

Restrictions: For Research Use only


Handling

Format: Lyophilized

Buffer: 50 mM Tris, 100 mM Glycine, 25 mM Arginine, 150 mM NaCl, pH 7.5


Storage: -20 °C

Images



ELISA

Image 1. Immobilized Human ACE2, His Tag (ABIN6952618, ABIN6952641) at 2 µg/mL (100 µL/well) can bind SARS-CoV-2 S1 protein, Fc Tag (ABIN6992404) with a linear range of 2-25 ng/mL (QC tested).



SDS-PAGE

Image 2. SARS-CoV-2 S1 protein, Fc Tag on under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95 % .