

Datasheet for ABIN6992411

IL2RG Protein (AA 23-254) (His tag)

1 Image



Go to Product page

Overview

Quantity:	100 μg
Target:	IL2RG
Protein Characteristics:	AA 23-254
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This IL2RG protein is labelled with His tag.

Product Details

Purpose:	Human IL-2 R gamma / CD132 Protein, His Tag (MALS verified)
Purity:	>95 % as determined by SDS-PAGE.
Endotoxin Level:	Less than 1.0 EU per µg by the LAL method.

Target Details

Target:	IL2RG
Alternative Name:	IL-2 R gamma (IL2RG Products)
Background:	IL-2R is a heterotrimeric protein binds and responds to the cytokine IL-2. Three distinct chains
	of IL-2R, termed as α , β and γ , which are non-covalently associated are identified. The α and β
	chains are involved in binding IL-2, while signal transduction following cytokine interaction is
	carried out by the γ chain, along with the β subunit. The α chain of the IL-2R can bind to the β
	chain before receptor interaction with IL-2. The γ chain alone has a very weak affinity for IL-2,

but after the ligand is bound to the α/β heterodimer, the γ chain becomes recruited to the complex to form a very stable macromolecular quaternary ligand/receptor complex. Interleukin-2 receptor subunit gamma (IL2RG), also known as cytokine receptor common subunit gamma, CD antigen CD132, gammaC, p64, which belongs to the type I cytokine receptor family or type 5 subfamily. IL2RG is located on the surface of immature blood-forming cells in bone marrow. Defects in IL2RG are the cause of severe combined immunodeficiency X-linked T-cell-negative/B-cell-positive/NK-cell-negative (XSCID).

Molecular Weight:

29.3 kDa

Pathways:

JAK-STAT Signaling, Growth Factor Binding

Application Details


Restrictions:

For Research Use only

Handling

Format:	Lyophilized
Buffer:	PBS, pH 7.4
Storage:	-20 °C

Images



SDS-PAGE

Image 1. Human IL-2 R gamma, His Tag on under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95 %.