antibodies

## Datasheet for ABIN700782 anti-RAPTOR antibody (AA 101-200) (Biotin)



Overview

Quantity:	100 µL
Target:	RAPTOR
Binding Specificity:	AA 101-200
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RAPTOR antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Raptor
lsotype:	lgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.
Target Details	
Target:	RAPTOR

Alternative N	Name:	Raptor (RAPTOR Products)
	Order at www.antiboo	ies-online.com   www.antikoerper-online.de   www.anticorps-enligne.fr   www.antibodies-online.cn

International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN700782 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

Background:	Synonyms: KOG1, Mip1, Regulatory-associated protein of mTOR, Raptor, p150 target of
	rapamycin (TOR)-scaffold protein, RPTOR, KIAA1303
	Background: Involved in the control of the mammalian target of rapamycin complex 1
	(mTORC1) activity which regulates cell growth and survival, and autophagy in response to
	nutrient and hormonal signals, functions as a scaffold for recruiting mTORC1 substrates.
	mTORC1 is activated in response to growth factors or amino acids. Growth factor-stimulated
	mTORC1 activation involves a AKT1-mediated phosphorylation of TSC1-TSC2, which leads to
	the activation of the RHEB GTPase that potently activates the protein kinase activity of
	mTORC1. Amino acid-signaling to mTORC1 requires its relocalization to the lysosomes
	mediated by the Ragulator complex and the Rag GTPases. Activated mTORC1 up-regulates
	protein synthesis by phosphorylating key regulators of mRNA translation and ribosome
	synthesis. mTORC1 phosphorylates EIF4EBP1 and releases it from inhibiting the elongation
	initiation factor 4E (eiF4E). mTORC1 phosphorylates and activates S6K1 at 'Thr-389', which
	then promotes protein synthesis by phosphorylating PDCD4 and targeting it for degradation.
	Involved in ciliogenesis.
Gene ID:	57521
UniProt:	Q8N122
Pathways:	Warburg Effect
Application Details	
Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 an
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should b

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN700782 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

## Handling

	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months