

Datasheet for ABIN700856 anti-NR1D1 antibody (Biotin)



_					
	1//	r	Vİ	\triangle	۸/
	V		VI		/ V

Quantity:	100 μL
Target:	NR1D1
Reactivity:	Human, Mouse, Rat, Cow
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NR1D1 antibody is conjugated to Biotin
Application:	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human NR1D1	
Isotype:	IgG	
Cross-Reactivity:	Cow, Human, Mouse, Rat	
Purification:	Purified by Protein A.	

Target Details

Target:	NR1D1	
Alternative Name:	NR1D1 (NR1D1 Products)	
Background:	Synonyms: EAR-1, EAR1, ERBA-RELATED 1, hRev, Nr1d1, NR1D1_HUMAN, Nuclear Receptor	
	Rev-ErbA Alpha, Nuclear receptor subfamily 1 group D member 1, Orphan nuclear receptor	
	NR1D1, Rev erbAalpha, Rev erbalpha, Rev-erbA-alpha, Rev-ErbAalpha, REV-ERBalpha, THRA1,	
	THRAL, Thyroid hormone receptor alpha-like, Thyroid hormone receptor, alpha like, V-erbA	

related protein EAR-1, V-erbA-related prote

Background: NR1D1, a NR1 Thyroid Hormone-Like Receptor, is encoded by the same genomic locus as, but transcribed from the opposite strand of, Thyroid Hormone Receptor Alpha (TR Alpha). NR1D1 is a target of Nuclear Receptor ROR Alpha and a transcription regulator that has been shown to affect myocyte differentiation, adipogenesis, and lipoprotein metabolism. Mice lacking NR1D1 show abnormal postnatal cerebellar development. NR1D1 expression has been documented in human skeletal muscle and a variety of mouse and rat tissues. ESTs have been isolated from human tissue libraries, including cancerous adrenal, blood, brain, breast, colon, duodenum, fetus, head/neck, kidney, lung, skeletal muscle, skin, synovium, uterus, normal brain, breast, colon, eye, heart, pancreas, pituitary, prostate, skeletal muscle, skin, testis and thyroid.

Pathways: Nuclear Receptor Transcription Pathway, Steroid Hormone Mediated Signaling Path	
Gene ID:	9572
Molecular Weight:	68kDa

Nuclear Receptor Transcription Pathway, Steroid Hormone Mediated Signaling Pathway,

Cellular Response to Molecule of Bacterial Origin, Regulation of Lipid Metabolism by PPARalpha

Application Details

WB(1:100-500)

Application Notes:

	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months