antibodies - online.com







anti-TRIM33 antibody (AA 981-1127) (Cy3)



| () | 1/0 | r\ /1 | 014 | |
|-----|-----|-------|-----|---|
| () | ve | I V I | -v | V |
| | | | | |

| Quantity: | 100 μL | |
|----------------------|--|--|
| Target: | TRIM33 | |
| Binding Specificity: | AA 981-1127 | |
| Reactivity: | Human, Mouse, Rat | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This TRIM33 antibody is conjugated to Cy3 | |
| Application: | Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) | |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human TIF1 gamma |
|-----------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Human, Mouse, Rat |
| Predicted Reactivity: | Dog,Cow,Pig,Horse,Chicken |
| Purification: | Purified by Protein A. |

Target Details

| Target: | TRIM33 |
|-------------------|----------------------------|
| Alternative Name: | T1 gamma (TRIM33 Products) |

Target Details

| Background: | Synonyms: ECTO, PTC7, RFG7, TF1G, TIF1G, TIFGAMMA, TIF1GAMMA, E3 ubiquitin-protein ligase TRIM33, Ectodermin homolog, RET-fused gene 7 protein, Protein Rfg7, Transcription intermediary factor 1-gamma, TIF1-gamma, Tripartite motif-containing protein 33, TRIM33, | |
|---------------------|--|--|
| | KIAA1113 | |
| | Background: Acts as an E3 ubiquitin-protein ligase. Promotes SMAD4 ubiquitination, nuclear exclusion and degradation via the ubiquitin proteasome pathway. According to | |
| | PubMed:16751102, does not promote a decrease in the level of endogenous SMAD4. May act | |
| | as a transcriptional repressor. Inhibits the transcriptional response to TGF-beta/BMP signaling | |
| | cascade. Plays a role in the control of cell proliferation. Its association with SMAD2 and SMAD3 | |
| | stimulates erythroid differentiation of hematopoietic stem/progenitor (By similarity). | |
| | Monoubiquitinates SMAD4 and acts as an inhibitor of SMAD4-dependent TGF-beta/BMP | |
| | signaling cascade (Monoubiquitination of SMAD4 hampers its ability to form a stable complex | |
| | with activated SMAD2/3 resulting in inhibition of TGF-beta/BMP signaling cascade). | |
| Gene ID: | 51592 | |
| UniProt: | Q9UPN9 | |
| Application Details | | |
| Application Notes: | FCM 1:20-100 | |
| | IF(IHC-P) 1:50-200 | |
| | IF(IHC-F) 1:50-200 | |
| | IF(ICC) 1:50-200 | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | |
| | handled by trained staff only. | |
| Storage: | -20 °C | |
| | | |

Handling

| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
|------------------|---|
| Expiry Date: | 12 months |