antibodies - online.com







anti-TRIM33 antibody (AA 981-1127) (Cy7)



()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL	
Target:	TRIM33	
Binding Specificity:	AA 981-1127	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This TRIM33 antibody is conjugated to Cy7	
Application:	Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human TIF1 gamma	
Isotype:	IgG	
Cross-Reactivity:	Human, Mouse, Rat	
Predicted Reactivity:	Dog,Cow,Pig,Horse,Chicken	
Purification:	Purified by Protein A.	

Target Details

Target:	TRIM33
Alternative Name:	T1 gamma (TRIM33 Products)

Target Details

Background:	Synonyms: ECTO, PTC7, RFG7, TF1G, TIF1G, TIFGAMMA, TIF1GAMMA, E3 ubiquitin-protein ligase TRIM33, Ectodermin homolog, RET-fused gene 7 protein, Protein Rfg7, Transcription intermediary factor 1-gamma, TIF1-gamma, Tripartite motif-containing protein 33, TRIM33, KIAA1113 Background: Acts as an E3 ubiquitin-protein ligase. Promotes SMAD4 ubiquitination, nuclear exclusion and degradation via the ubiquitin proteasome pathway. According to PubMed:16751102, does not promote a decrease in the level of endogenous SMAD4. May act as a transcriptional repressor. Inhibits the transcriptional response to TGF-beta/BMP signaling	
	cascade. Plays a role in the control of cell proliferation. Its association with SMAD2 and SMAD3	
	stimulates erythroid differentiation of hematopoietic stem/progenitor (By similarity).	
	Monoubiquitinates SMAD4 and acts as an inhibitor of SMAD4-dependent TGF-beta/BMP	
	signaling cascade (Monoubiquitination of SMAD4 hampers its ability to form a stable complex	
	with activated SMAD2/3 resulting in inhibition of TGF-beta/BMP signaling cascade).	
Gene ID:	51592	
UniProt:	Q9UPN9	
Application Details		
Application Notes:	FCM 1:20-100	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	

Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months