antibodies .- online.com





Datasheet for ABIN701147

anti-TRIM33 antibody (AA 981-1127) (FITC)



Overview

Quantity:	100 μL
Target:	TRIM33
Binding Specificity:	AA 981-1127
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TRIM33 antibody is conjugated to FITC
Application:	Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human TIF1 gamma
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Pig,Horse,Chicken
Purification:	Purified by Protein A.

Target Details

Target:	TRIM33
Alternative Name:	T1 gamma (TRIM33 Products)

Target Details

Background:	Synonyms: ECTO, PTC7, RFG7, TF1G, TIF1G, TIFGAMMA, TIF1GAMMA, E3 ubiquitin-protein
	ligase TRIM33, Ectodermin homolog, RET-fused gene 7 protein, Protein Rfg7, Transcription
	intermediary factor 1-gamma, TIF1-gamma, Tripartite motif-containing protein 33, TRIM33,
	KIAA1113
	Background: Acts as an E3 ubiquitin-protein ligase. Promotes SMAD4 ubiquitination, nuclear
	exclusion and degradation via the ubiquitin proteasome pathway. According to
	PubMed:16751102, does not promote a decrease in the level of endogenous SMAD4. May act
	as a transcriptional repressor. Inhibits the transcriptional response to TGF-beta/BMP signaling
	cascade. Plays a role in the control of cell proliferation. Its association with SMAD2 and SMAD3
	stimulates erythroid differentiation of hematopoietic stem/progenitor (By similarity).
	Monoubiquitinates SMAD4 and acts as an inhibitor of SMAD4-dependent TGF-beta/BMP
	signaling cascade (Monoubiquitination of SMAD4 hampers its ability to form a stable complex
	with activated SMAD2/3 resulting in inhibition of TGF-beta/BMP signaling cascade).
Gene ID:	51592
UniProt:	Q9UPN9
Application Details	
Application Notes:	FCM 1:20-100
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C

Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months