

### Datasheet for ABIN701209

# anti-Tuberin antibody (AA 1601-1750) (HRP)



Go to Product page

_					
	1//	r	Vİ	$\triangle$	۸/
	V		VI		/ V

Quantity:	100 μL	
Target:	Tuberin (TSC2)	
Binding Specificity:	AA 1601-1750	
Reactivity:	Human, Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Tuberin antibody is conjugated to HRP	
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human Tuberin	
Isotype:	IgG	

## Cross-Reactivity:

Human, Mouse, Rat

#### Predicted Reactivity:

Dog,Cow,Pig,Horse,Rabbit

#### Purification:

Purified by Protein A.

## **Target Details**

Target: Tuberin (TSC2)

Alternative Name: Tuberin (TSC2 Products)

## **Target Details**

J				
Background:	Synonyms: LAM, TSC4, PPP1R160, Tuberin, Tuberous sclerosis 2 protein, TSC2			
	Background: In complex with TSC1, inhibits the nutrient-mediated or growth factor-stimulated			
	phosphorylation of S6K1 and EIF4EBP1 by negatively regulating mTORC1 signaling. Acts as a			
	GTPase-activating protein (GAP) for the small GTPase RHEB, a direct activator of the protein kinase activity of mTORC1. Implicated as a tumor suppressor. Involved in microtubule-mediated protein transport, but this seems to be due to unregulated mTOR signaling.  Stimulates weakly the intrinsic GTPase activity of the Ras-related proteins RAP1A and RAB5 in			
	vitro. Mutations in TSC2 lead to constitutive activation of RAP1A in tumors.			
Gene ID:	7249			
UniProt:	P49815			
Pathways:	RTK Signaling, AMPK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling			
	Pathway, Neurotrophin Signaling Pathway, Regulation of Cell Size, Tube Formation, Protein			
	targeting to Nucleus			
Application Details				
Application Notes:	IHC-P 1:200-400			
	IHC-F 1:100-500			
Restrictions:	For Research Use only			
Handling				
Format:	Liquid			
Concentration:	1 μg/μL			
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.			
Preservative:	ProClin			
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.			
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.			
Storage:	-20 °C			
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.			

		1.	
$\vdash$	land	ling	
1 1	ıaııu	11111	۰

Expiry Date:

12 months