# antibodies -online.com





# CD56 Protein (AA 20-718) (Fc Tag)

2 Images



Go to Product page

#### Overview

Quantity:	100 μg
Target:	CD56 (NCAM1)
Protein Characteristics:	AA 20-718
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This CD56 protein is labelled with Fc Tag.

### **Product Details**

Characteristics:	Human NCAM-1 / CD56 Protein, Fc Tag (MALS verified)
Purity:	>95 % as determined by SDS-PAGE.
Endotoxin Level:	Less than 1.0 EU per μg by the LAL method.

# **Target Details**

Target:	CD56 (NCAM1)
Alternative Name:	NCAM-1 (NCAM1 Products)
Background:	NCAM1 belongs to the immunoglobulin superfamily of adhesion molecules. A wide range of alternatively spliced NCAM1 messenger RNAs (mRNAs) has been described to date, but only the 120-, 140-, and 180- kDa isoforms are commonly expressed. NCAM1 plays an important role in the regulation of neurogenesis, neurite outgrowth, proliferation, and cell migration,

however, its function in hematopoiesis, including NK cells, is poorly understood. NCAM1 signaling is mediated either by homophilic or heterophilic interactions with fibroblast growth factor receptor (FGFR), L1-CAM, N-cadherin and other components of the extracellular matrix. Upon activation, NCAM1 triggers a variety of signaling cascades including FYN-focal adhesion kinase (FAK), MAPK, and phosphatidylinositol 3-kinase (PI3K) pathways.

Molecular Weight: 104.0 kDa

NCBI Accession: NP\_851996


## **Application Details**

Application Notes:	MALS verified
Restrictions:	For Research Use only

### Handling


Format:	Lyophilized
Buffer:	PBS, pH 7.4
Storage:	-20 °C

#### **Images**



#### **ELISA**

**Image 1.** Immobilized Human NCAM-1, Fc Tag (ABIN6973166) at  $1 \mu g/mL$  (100  $\mu L/well$ ) can bind A-1 Antibody, Human IgG1 with a linear range of 0.3-10 ng/mL (QC tested).



### **SDS-PAGE**