antibodies -online.com





CD56 Protein (AA 20-718) (Fc Tag,FITC)

2 Images



Go to Product page

Overview

Quantity:	200 μg
Target:	CD56 (NCAM1)
Protein Characteristics:	AA 20-718
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This CD56 protein is labelled with Fc Tag,FITC.

Product Details

Purpose:	FITC-Labeled Human NCAM-1 / CD56 Protein, Fc Tag
Purity:	>90 % as determined by SDS-PAGE.
Endotoxin Level:	Less than 1.0 EU per µg by the LAL method.

Target Details

Target:	CD56 (NCAM1)
Alternative Name:	NCAM-1 (NCAM1 Products)
Background:	NCAM1 belongs to the immunoglobulin superfamily of adhesion molecules. A wide range of
	alternatively spliced NCAM1 messenger RNAs (mRNAs) has been described to date, but only
	the 120-, 140-, and 180- kDa isoforms are commonly expressed. NCAM1 plays an important
	role in the regulation of neurogenesis, neurite outgrowth, proliferation, and cell migration,

however, its function in hematopoiesis, including NK cells, is poorly understood. NCAM1 signaling is mediated either by homophilic or heterophilic interactions with fibroblast growth factor receptor (FGFR), L1-CAM, N-cadherin and other components of the extracellular matrix. Upon activation, NCAM1 triggers a variety of signaling cascades including FYN-focal adhesion kinase (FAK), MAPK, and phosphatidylinositol 3-kinase (PI3K) pathways.

Molecular Weight: 104.0 kDa

NCBI Accession: NP_851996


Application Details

Restrictions: For Research Use only

Handling


Format:	Lyophilized
Buffer:	PBS, pH 7.4
Storage:	-20 °C

Images



ELISA

Image 1. Immobilized A-1 Antibody, Human IgG1 at $5\,\mu$ g/mL (100 μ L/well) can bind Fed Human NCAM-1, Fc Tag (ABIN6992429) with a linear range of 0.08-0.625 μ g/mL (QC tested).



SDS-PAGE

Image 2. Fed Human NCAM-1, Fc Tag on under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 90 %.