antibodies -online.com





TNFRSF13C Protein (AA 7-71) (His tag)

3 In

Images



Go to Product page

()	1 /	-	rv	1 /	71	A
	1//	\vdash	1 \/	16		۱/۱
\sim	٧.	\sim	1 V	ı١	_	V١

Quantity:	100 μg
Target:	TNFRSF13C
Protein Characteristics:	AA 7-71
Origin:	Human
Source:	HEK-293 Cells
Protein Type:	Recombinant
Biological Activity:	Active
Purification tag / Conjugate:	This TNFRSF13C protein is labelled with His tag.

Product Details

Characteristics:	Human BAFFR / TNFRSF13C Protein, His Tag
Purity:	>90 % as determined by SDS-PAGE.
Endotoxin Level:	Less than 1.0 EU per μg by the LAL method.

Target Details

Target:	TNFRSF13C
Alternative Name:	BAFFR (TNFRSF13C Products)
Background:	BAFF receptor (B-cell activating factor receptor, BAFF-R), also known as tumor necrosis factor receptor superfamily member 13C (TNFRSF13C), is a membrane protein of the TNF receptor superfamily which recognizes BAFF. B-cell activating factor (BAFF) enhances B-cell survival in vitro and is a regulator of the peripheral B-cell population. Overexpression of BAFF in mice

results in mature B-cell hyperplasia and symptoms of systemic lupus erythematosus (SLE). Also, some SLE patients have increased levels of BAFF in serum. Therefore, it has been proposed that abnormally high levels of BAFF may contribute to the pathogenesis of autoimmune diseases by enhancing the survival of autoreactive B cells.

Molecular Weight:	8.6 kDa
NCBI Accession:	NP_443177
Pathways:	NF-kappaB Signaling


Application Details

Restrictions: For Research Use only

Handling


Format:	Lyophilized	
Buffer:	PBS, pH 7.4	
Storage:	-20 °C	

Images




ELISA

Image 1. Immobilized Human BAFFR Protein, His Tag (ABIN6972953) at $2\,\mu g/mL$ (100 $\mu L/well$) can bind A Antibody, Human IgG1 with a linear range of 0.1-5 ng/mL (Routinely tested).



SDS-PAGE

Image 2. Human BAFFR Protein, His Tag on under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 90 %.



ELISA

Image 3. Immobilized Human BAFF, Fc Tag, active trimer (ABIN6972950) at 1 μ g/mL (100 μ L/well) can bind Human BAFFR Protein, His Tag (ABIN6972953) with a linear range of 1-10 ng/mL (QC tested).