

Datasheet for ABIN7013920

anti-HLA-E antibody**2** Images[Go to Product page](#)

Overview

Quantity:	0.1 mg
Target:	HLA-E
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This HLA-E antibody is un-conjugated
Application:	Flow Cytometry (FACS), ELISA, Immunoprecipitation (IP)

Product Details

Immunogen:	recombinant human HLA-E
Clone:	3D12
Isotype:	IgG1
Specificity:	The mouse monoclonal antibody 3D12 (also known as 3D12HLA-E) recognizes native extracellular part of HLA-E, an ubiquitously expressed non-classical MHC class I molecule, as well as free HLA-E.
Purification:	Purified by protein-A affinity chromatography.

Target Details

Target:	HLA-E
Alternative Name:	HLA-E (HLA-E Products)

Target Details


Background:	Major histocompatibility complex, class I, E,HLA-E (human leukocyte antigen E) is a non-classical MHC I antigen, which is important for dialogue with NK cells and their regulation through interaction with CD94/NKG2 receptor. Like other MHC I molecules, transmembrane HLA-E molecule (45 kDa) associates with beta2 microglobulin. Unlike HLA-G, expression of HLA-E molecules is not so restricted, but it has been detected at least at mRNA level in virtually all cells and tissues examined. In peripheral blood, HLA-E protein is expressed at least in all mononuclear cells, but in different quantity (B cells and monocytes more than T cells and NK cells).,HLA class I histocompatibility antigen, alpha chain E, MHC class I antigen E
Gene ID:	3133
UniProt:	P13747
Pathways:	Regulation of Leukocyte Mediated Immunity , Positive Regulation of Immune Effector Process

Application Details

Application Notes:	Flow cytometry: Recommended dilution: 1-4 µg/mL.
Restrictions:	For Research Use only

Handling

Concentration:	1 mg/mL
Buffer:	Phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Store at 2-8°C. Do not freeze.



Flow Cytometry

Image 1. Flow cytometry surface staining pattern of human peripheral blood stained using anti-human HLA-E (3D12) purified antibody (concentration in sample 4 µg/mL) GAM APC.

Flow Cytometry

Image 2. Separation of human lymphocytes (red-filled) from blood debris (black-dashed) in flow cytometry analysis (surface staining) of human peripheral whole blood stained using anti-human HLA-E (3D12) purified antibody (concentration in sample 4 µg/mL) GAM APC.